


# Controlling pressure-driven ionic flow by voltage at molecular scale

March 7 2019, by Charlotte Powell

---



Credit: University of Manchester

Similar to our computers which handle electrons to perform the calculations and logics, all the circuitry in living beings is based on the transport of ions, such as sodium, chloride, calcium, etc. Nature exploits incredibly subtle transport of these elementary charges and an artillery of ion channels to perform advanced functions by manipulating the—often

exotic—behaviour of ion transport at molecular scales. Achieving such features in artificial channels remains a considerable challenge.

As published in *Nature*, Researchers from the Micromegas team at the Physics Department at ENS, Paris in collaboration with the Condensed Matter Physics laboratory and National Graphene Institute at The University of Manchester, have been able to highlight mechano-sensitive properties of ion [transport](#) in few angstroms thick artificial channels.

Just over two years ago, Manchester researchers led by Dr. Radha Boya and Prof Sir Andre Geim showed that by stacking two-dimensional atomic layers similar to stacking bricks of Lego, it is indeed possible to assemble molecular and smooth channels at the atomic scale in a controlled manner. The atomic layers used for building the [channel](#) are held together by so-called van der Waals forces. Using these channels, the new experiments show that considerable ionic current can be generated when a flow is induced by applying a pressure difference. Separating two miniature baths of salt solutions, these angstrom scale channels generate ionic current when water molecules are mechanically pushed through them.

Dr. Timothée Mouterde, the first author of this study, said: "Even more surprising, by applying an [electric field](#) along with pressure, this flow current can be modulated extremely sensitively."

Prof Lydéric Bocquet adds: "This novel effect is akin to transistor but here for ion transport and can be understood as gating of mechanical ion [flow](#) by voltage." Furthermore interestingly, the electronic properties of the confining wall materials of the channel seem to influence this 'voltage gating.' This effect can be understood by differential friction of water and ions on the walls at these molecular scales."

Dr. Ashok Keerthi, who is a co-author said: "Inside our artificial

channels which are only couple of water atoms thick, water and ions are organized in a two-dimensional monolayer. The ability to make such precise angstrom scale channels has provided us with tools to explore anomalous properties of [water](#) and flows."

Dr. Radha Boya explains: "At the molecular [scale](#), flows induced by pressure and voltage simply do not add up. This coupling between mechanical and electrical forces demonstrated at the ultimate scales shows strong similarities to those observed in mechanically sensitive biological ion channels such as PIEZO1. This new platform will allow exploring the physical mechanisms of these extreme confinement situations at work in living systems, and in the longer term, to mimic elementary calculus functions based on ion transport."

**More information:** A. Keerthi et al. Ballistic molecular transport through two-dimensional channels, *Nature* (2018). [DOI: 10.1038/s41586-018-0203-2](#)

Provided by University of Manchester

Citation: Controlling pressure-driven ionic flow by voltage at molecular scale (2019, March 7) retrieved 20 April 2024 from <https://phys.org/news/2019-03-pressure-driven-ionic-voltage-molecular-scale.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.