

Probing water's skin

March 7 2019



KAUST researchers have used complementary techniques to disentangle purely interfacial effects from electrospray-specific effects. Credit: Ivan Gromicho

From the wind-whipped surface of the open ocean, to trillions of tiny water drops in clouds, the air-water interface—water's skin—is the site for crucial natural processes, including ocean-atmosphere exchange and cloud acidification. The air-water interface has even been postulated as the cradle of life. However, factors such as its subnanometer size and dynamic nature, render the investigation of interfacial water a daunting task.

In recent years, researchers have investigated the air-water [interface](#) using electrosprays of water: fine sprays produced by applying greater than 5000 V to water solutions passing through a metallic capillary. Traditionally, electrosprays have been used to study ions in the gas phase. Using electrosprays, researchers have suggested that the surface of mildly acidic water behaves as a highly reactive superacid. But the debate remains whether the air-water interface at mild pH can really behave as a superacid.

Now, KAUST researchers led by Himanshu Mishra have employed complementary techniques to disentangle purely interfacial effects from [electrospray](#)-specific effects.


The researchers investigated the reactivity of isoprene—a volatile molecule released by plants experiencing heat stress—at the water interface. "We compared two scenarios: electrosprays of water interacting with isoprene gas and vigorously stirred mixtures of water and isoprene in closed vials," explains Adair Gallo Jr., a Ph.D. student in Mishra's team and lead author of the study.

The team looked for the formation of short isoprene chains called oligomers. "Intriguingly, isoprene spontaneously formed oligomers in electrosprays, under acidic and basic conditions, and even in the absence of water," Gallo says. No oligomerization products were detected when acidified water was vigorously stirred with [isoprene](#) for hours. But when the same organic phase was electrosprayed, oligomers formed. The findings unequivocally establish that the oligomerization took place exclusively in electrosprays.

Computer simulations carried out by Adriano Sanchez, a postdoctoral scholar in Mishra's team, gave molecular-scale insights into the results. "We found that oligomer formation was only possible on gas-phase clusters comprising not more than three water molecules and an excess

proton," Sanchez says.

Collectively, the team's results demonstrated that electrosprays present highly energetic [gas-phase](#) pathways for [chemical reactions](#) to occur that are impossible at natural air-[water](#) interfaces. "Electrosprays should therefore be complemented with surface-specific techniques and [computer simulations](#) to avoid incorrect conclusions when studying interfacial processes," Mishra says. "I have been thinking about this problem for over six years, and now, thanks to my team, we have managed to disentangle purely interfacial effects from artifacts" says Mishra. This contribution will feature on the cover of the journal *Chemical Science*.



Electrospray ionization can produce tiny highly charged and acidic droplets that

catalyze chemical reactions in the gas phase. Credit: Ivan Gromicho

More information: Adair Gallo et al. The chemical reactions in electrosprays of water do not always correspond to those at the pristine air–water interface, *Chemical Science* (2018). [DOI: 10.1039/C8SC05538F](https://doi.org/10.1039/C8SC05538F)

Provided by King Abdullah University of Science and Technology

Citation: Probing water's skin (2019, March 7) retrieved 23 May 2024 from <https://phys.org/news/2019-03-probing-skin.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.