


# Small brains, big picture: Study unveils *C. elegans*' microscopic mysteries

March 5 2019

---



(Left to right) Dr. Yuto Momohara, Dr. Takashi Murayama, Prof. Ichiro Maruyama, Professor Jeff Wickens and Dr. Ei-ichiro Saita started a collaboration between the Information Processing Biology Unit and the Neurobiology Research Unit to study the neurons in the *C. elegans* brain. Not pictured are first author Dr. Tomomi Shindou and Dr. Mayumi Ochi-Shindou.  
Credit: OIST

It's the small pieces that make the big picture, and in this case, the pieces can't be seen by the naked eye. New research at the Okinawa Institute of Science and Technology Graduate University (OIST) used microscopy techniques to piece together the brain of the millimeter-long *Caenorhabditis elegans*, revealing that their neurons fire action potentials—a spike in voltage when neurons transmit sensory information in the cell membrane. Their results could lead to better understanding of nerve signal transmission in the organism and serve as a future model for neuronal information processing in other animals, including humans.

Scientists from the Information Processing Biology Unit (IPBU) and the Neurobiology Research Unit (NRU) at OIST collaborated on the project. The IPBU had the technology to genetically modify *C. elegans*, while the NRU had experience recording electrical signals from the worm's tiny [neurons](#). Although generally thought to lack the ability to fire [action potentials](#)—the all-or-none pulses needed for decision-making—Prof. Ichiro Maruyama, principal investigator of the IPBU, had results that suggested the worm's neurons did fire action potentials. To confirm his suspicions, he wanted further data using electrophysiology, which his lab was not actively doing, so he shared his ideas with Prof. Jeff Wickens, head of the NRU and researchers Tomomi and Mayumi Shindou.

Seven years later, the collaboration paid off; the completed study was published in *Scientific Reports* on March 5, 2019. Before the project, the common perception was that *C. elegans* neurons only responded in a graded way with analog signals. The work showed, however, that the neurons would fire an action potential if they crossed a threshold, a more digital type of signaling.

"The main thing this paper shows is that the neurons we study fire action

potentials," Wickens said. "Now, *C. elegans* neurons have both active and passive transmission, so all these mechanisms are essential for sensory processing, even in the smallest of neurons."

To observe sensory processing in *C. elegans*, Maruyama and his team applied salt to the tip of the worm's nose. They increased and decreased the salt concentration, seeking different reactions from the worm and monitoring the electroactivity of its neurons. They found that an increase in salt first affected a neuron on the left side of the brain, then triggered signals causing the worm to move forward. A decrease caused a neuron on the right side to send signals for the worm to move backwards.

"This pair of neurons have the same structure and name, but completely opposite functions," Maruyama said. "Using those two neurons together is very efficient for the worm to reach a place with a preferable salt concentration."

## **Developing Miniscule Methods**

Researching *C. elegans* imposes a challenge—the brain of *C. elegans* has 302 neurons, while an ant has 250,000 and a mouse has 75 million. Wickens put his former postdoc and first author of the paper, Dr. Tomomi Shindou, in charge of developing the methodology for the project. The work took four years of trial and error to create the right techniques, and three more for collecting data.

The NRU created a sharp glass needle less than the size of a micrometer. They used the needle to extract a brain cell from *C. elegans* after the worm was exposed to salt. This method is like using tweezers to extract a splinter from a finger.

"Only a handful of labs in the world can do this," Maruyama said. "Once the cell was extracted, the scientists used electrophysiology—measuring

voltage change in cells to track neurons viewing active and passive activation. The scientists also used calcium imaging to produce more data, and observed freely moving worms that had been exposed to the salt.

"You can express a calcium sensor protein in a particular neuron—if the neuron is activated, it will light up," Maruyama said. "Once neurons are activated, they produce a green fluorescence, and now you are detecting the neuron activity with the fluorescent microscope. The data from the two tests perfectly matched."

"Perfectly matched data" is a phrase every researcher hopes to hear. The results suggest that the accepted dogma—that these worms only send passive signals—is wrong. The remarkable finding also opens a door to improved neural mapping in the future.

"Every *C. elegans* has the same set of neurons connected in the same way, unlike a mouse brain, which is a jungle of connections," Wickens said. "If we can get a better understanding of this simple nervous system, for which we have a kind of circuit diagram, we can get insights on how more complicated brains operate at a bigger scale."

**More information:** Tomomi Shindou et al. Active propagation of dendritic electrical signals in *C. elegans*, *Scientific Reports* (2019). [DOI: 10.1038/s41598-019-40158-9](https://doi.org/10.1038/s41598-019-40158-9)

Provided by Okinawa Institute of Science and Technology

Citation: Small brains, big picture: Study unveils *C. elegans*' microscopic mysteries (2019, March 5) retrieved 23 April 2024 from <https://phys.org/news/2019-03-small-brains-big-picture-unveils.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.