

Tasty or putrid? A repellent odor inhibits the perception of a pleasant odor in vinegar flies

March 18 2019



Vinegar flies process conflicting odors in the brain by impairing the perception of good odors through neuronal interaction with noxious odors. Credit: Benjamin Fabian

In nature, vinegar flies are exposed to a wide variety of odor mixtures,

which contain both attractive and repellent odors. Scientists at the Max Planck Institute for Chemical Ecology have now discovered that repellent odors suppress the perception of pleasant smells. This happens because certain brain structures that respond to attractive odors are inhibited by a repellent one. These processes in the brain are also reflected in the behavior of the flies: The flies avoid attractive odor sources when a repellent odor is added. This helps them to avoid spoiled or infected food sources, which would have fatal consequences for the flies and their offspring.

Vinegar flies are always exposed to odor mixtures when navigating in the natural environment. These mixtures often consist of odors which are both attractive and repellent, such as a source of food contaminated with toxic bacteria. The decision to approach or avoid is extremely important for the survival and reproduction of the animal. But so far, little is known about how different odors with opposing values in a mixture are processed simultaneously in the brain. In previous studies, the research groups led by Silke Sachse and Markus Knaden from the Max Planck Institute for Chemical Ecology have already shown that the olfactory channels that respond to attractive scents are different from those which are activated by repellent odors. In addition, two pleasant odors reinforce each other in certain [neural circuits](#) in the fly brain, which is in turn reflected in behavior.

"In this study, we aimed to investigate how odor mixtures of opposite valence, that is, a mix of odors that are attractive and repellent, are processed and perceived," explains Ahmed Mohamed, lead author of the study, adding: "By taking advantage of the sophisticated genetics available in *Drosophila*, we elucidated the neural mechanism that enables the animal to evaluate such conflicting situations in order to take the appropriate decision."

The scientists exposed the vinegar fly *Drosophila melanogaster* to

artificial odor mixtures, each containing an attractive and repellent odor, in defined ratios. By analyzing the [brain activity](#) of these opposing odor mixtures using functional imaging techniques, they were able to show that a repellent component in an odor mixture specifically inhibits the odor channels for attractants. "This inhibition correlates with a reduced attraction to the odor source" explains Silke Sachse. The underlying neuronal mechanism and the specific neurons involved could also be identified.

"Our data demonstrate a specific, inhibitory cross-talk in the olfactory center of the fly's brain. Using further genetic tools we were able to show that glomeruli, spherical functional units in the olfactory center, that respond to attractive odors, are linked via specific inhibitory neurons to the repellent-specific glomeruli," Ahmed Mohamed describes from the results.

However, not all [odor](#) mixtures show the same inhibitory effect. An exception is [geosmin](#), the typical smell of toxic bacteria or mold. Geosmin is recognized in the fly brain by only one receptor type, and thus only a highly specific glomerulus is activated. "It is conceivable that this geosmin glomerulus has no strong interactions with other glomeruli and cannot influence them accordingly," speculates Markus Knaden, another author of the study. Similar specialized pathways have been described only for the detection of sex pheromones, carbon dioxide and iridomyrmecin, the latter of which is the smell specific to a parasitic wasp that infests [vinegar flies](#).

The scientists assume that this neuronal mechanisms of the fly [brain](#) might also occur in other olfactory systems, such as the olfactory bulb of mammals, and therefore also potentially in humans. Such a mechanism could help protect not only flies, but also we as humans to recognize and avoid, for example, contaminated food that emits pleasant as well as bad odors. In the future, the researchers want to investigate whether this

neuronal mechanism is modified by other factors, such as physiological state, e.g. hunger.

More information: Ahmed A. M. Mohamed et al. Odor mixtures of opposing valence unveil inter-glomerular crosstalk in the *Drosophila* antennal lobe, *Nature Communications* (2019). [DOI: 10.1038/s41467-019-09069-1](https://doi.org/10.1038/s41467-019-09069-1)

Provided by Max Planck Society

Citation: Tasty or putrid? A repellent odor inhibits the perception of a pleasant odor in vinegar flies (2019, March 18) retrieved 13 March 2024 from <https://phys.org/news/2019-03-tasty-putrid-repellent-odor-inhibits.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--