

Temperature blob in Pacific Ocean gives a glimpse of climate impact on humpback whales

March 20 2019, by Bob Yirka



Credit: CC0 Public Domain

A team of researchers with the Keiki Kohola Project and California State University reports that a recent "blob" of warm water in the Pacific



gives marine scientists a preview of the impact of climate change on humpback whales. In their paper published in the journal *Royal Society Open Science*, the group describes their study of the whales and what they found.

Humpback whales were once hunted to the point that they became endangered, but several decades ago, <u>international laws</u> were enacted to protect them, and they rebounded—these measures were so successful that the U.S. National Marine Fisheries Service took the whales off the <u>endangered species list</u> back in 2016. But now, it appears the whales face a new threat—climate change.

In a new effort, the researchers studied the reproductive habits of humpback whales that migrate between the waters off the Alaskan coast and those that surround the Hawaiian Islands. They had noted that in 2013, a giant mass or "blob" of warm <u>water</u> had formed in the Gulf of Alaska, and over the course of the next six years, slowly made its way south. Prior research had shown that temperatures in the blob were up to 3 degrees C warmer than surrounding water—high enough to kill krill and other sea creatures that serve as food for humpback whales. This led the researchers to embark on a study of the impact of the blob on the humpback whales.

According to prior research, when female <u>humpback</u> whales have insufficient food to pack on blubber, they stop ovulating. And if ovulation does not occur, a female will skip the migration season. In studying the whales, the researchers found that fewer females migrated when the water was warm, which meant that fewer <u>whales</u> were born. They were not able to pinpoint the actual number of births, but report that over the years 2013 and 2014, they counted one mother and calf pair for every three kilometers of ocean. In contrast, they had to search over 12 kilometers of ocean in 2017 and 2018 to find a mother/calf pair.



The researchers report that the blob has finally dissipated, but point out that its presence provides a glimpse of what <u>humpback whales</u> could be facing if the ocean warms as predicted in the coming years due to climate change.

More information: R. Cartwright et al. Fluctuating reproductive rates in Hawaii's humpback whales, Megaptera novaeangliae, reflect recent climate anomalies in the North Pacific, *Royal Society Open Science* (2019). DOI: 10.1098/rsos.181463

© 2019 Science X Network

Citation: Temperature blob in Pacific Ocean gives a glimpse of climate impact on humpback whales (2019, March 20) retrieved 3 May 2024 from <u>https://phys.org/news/2019-03-temperature-blob-pacific-ocean-glimpse.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.