


Tiny 'blinkers' enable simultaneous imaging of multiple biomolecules

March 13 2019

Protein barcoding



To image two proteins (CHC and PMP70) with the same fluorophore (green star), researchers targeted the proteins with two different lengths of DNA sequences attached to antibodies (Y-shaped molecules). Credit: American Chemical Society

In everyday life, blinking lights can send signals—for example, that a car is going to turn. Now, researchers have engineered tiny "blinkers" that reveal single molecules of RNA or protein inside cells based on the duration and frequency of each flash. The research, published in the ACS journal *Nano Letters*, could allow scientists to see the locations of many different biomolecules in a cell simultaneously, possibly leading to better diagnostics and treatments.

Recently, scientists have developed super-resolution microscopes that can image single [molecules](#) that are only a few nanometers in size. To discriminate a specific nucleic acid or protein, they typically add a [fluorescent probe](#) that binds to that molecule and emits a certain wavelength of light. However, because the emission wavelengths of different fluorescent probes can overlap, researchers can usually only detect three or four unique proteins or [nucleic acids](#) at a time, instead of the thousands that exist in cells. Ralf Jungmann and colleagues wondered if they could use fluorescent probes that blink with light at a variable duration and frequency to detect dozens of biomolecules at once. That way, they could use a single fluorophore to image many different molecules.

The researchers based their system on complementary sequences of DNA that come together to link a fluorophore with a target biomolecule and then fall apart again, generating a blinking fluorescent signal. By varying the length and number of DNA sequences bound to the target, the researchers could adjust how long the blink lasted, as well as how often blinks occurred. To test their approach in [cells](#), the researchers imaged two different RNA molecules and two proteins. Then, they used three fluorescent probes to image 124 distinct DNA structures that contained different numbers of target DNAs so that they blinked at different frequencies. The procedure took only a few minutes and had an accuracy of 97.6 percent, the researchers say.

More information: "124-Color Super-resolution Imaging by Engineering DNA-PAINT Blinking Kinetics" *Nano Letters* (2019).
pubs.acs.org/doi/abs/10.1021/acs.nanolett.9b00508

Provided by American Chemical Society

Citation: Tiny 'blinkers' enable simultaneous imaging of multiple biomolecules (2019, March 13)
retrieved 9 April 2024 from
<https://phys.org/news/2019-03-tiny-blinkers-enable-simultaneous-imaging.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--