

Trade-offs in adaptive anatomy: Vinegar fly species have good vision or olfaction, but not both

March 14 2019



A comparison of 62 different genera of *Drosophila* flies revealed that vinegar flies have either developed a more sensitive odor perception at the cost of poor vision or vice versa. Credit: Anna Schroll

A team of scientists from the Max Planck Institute for Chemical Ecology has systematically studied and compared the eyes and antennae and the associated brain structures of more than 60 species of the genus *Drosophila*. They found a large variation, but at the same time a close correlation between the two sensory features. The size of the sensory organs is related to the behavior of the mate and host selection.

The authors had observed different behavior in earlier studies on the black-bellied vinegar fly *Drosophila melanogaster* and the cherry vinegar fly *Drosophila suzukii*, a relatively new pest in Central Europe: while *D. melanogaster* are more likely to be attracted by the smell of food alone, in *D. suzukii*, also relies on vision in foraging. These sensory preferences were also reflected in the different sizes of the respective sensory organs.

Based on this observation, the scientists hypothesized that the variation in vision and olfaction is the result of a trade-off, in which a beneficial change in one trait is linked to a detrimental change in another trait in adaptation to the environment. In the case of the genus *Drosophila*, there is an apparent trade-off in the expression of visual and olfactory organs.

In order to test their trade-off hypothesis, the researchers examined the forms and functions of eyes and antennae as well as the associated visual and olfactory brain structures of a total of 62 *Drosophila* species: "The genus *Drosophila* offered us a much greater variation in sensory expression than we anticipated within a single closely related group of insects. And indeed, big-eyed species had small noses or antennae, while species with larger antennae had proportionally smaller eyes," says lead author Ian Keesey. The size of the sense organs, in turn, reflects the preferences related to selecting a host plant or a mate. Big-nose flies are more likely to use olfactory cues, while big-eyed flies are more likely to follow visual cues.

"The detailed analysis of eyes and antennae revealed the whole spectrum of the trade-off between vision and olfaction: We found species that had invested primarily in vision, species where vision and olfaction are about equal, and species that rely primarily on their olfactory sense, but none of the species studied had both large eyes and large antennae," explains Markus Knaden. For their analyses, the researchers reconstructed the primary sensory brain structures that play a role in vision and olfaction, including the optic and antennal lobes.

In addition, they used high-resolution microscopy to take a closer look at the sensory organs of the various fly species. "One reason why animals have to choose either a well-developed olfactory system or vision might be that in embryonic development both sensory organs emerge from the same structure with only a limited number of nuclei. The competition for resources, which decides which of the two sensory organs is more pronounced, thus takes place at a very early stage of development," says Bill Hansson, head of the Department of Evolutionary Neuroethology.

An important finding of the study is that genetic traits are linked. A change in one trait can have a large impact on the organism. However, some properties are less easy to modify, especially when tethered together with another. "It is fascinating that two senses so well studied, such as [vision](#) and olfaction, are inversely correlated. We now suspect that evolutionary pressures exist that are driving insects to prioritize the eye or the nose," says Ian Keesey.

With their study, the scientists want to open new avenues in the so-called Eco-Evo-Devo research. This [research field](#) is based on the assumption that concepts of ecology (eco), evolution (evo) and developmental biology (devo) are tightly linked, and the understanding of ecological relationships also requires evolutionary and developmental knowledge and vice versa. Although genomic data are available for many species, knowledge of their ecology is often lacking. "These trade-offs, especially

in genetic model organisms, provide an avenue for determining the mechanisms for how ecology and evolution shape the natural world," says Ian Keeseey.

The scientists also want to encourage other research groups not to look only at the well-known *Drosophila melanogaster*, but to include more species of this genus in their studies. The extensive datasets of this study are all available in the Max Planck database [Edmond](#) and researchers are welcome to use them for comparisons with other *Drosophila* [species](#) of the entire genus.

More information: Ian W. Keeseey et al, Inverse resource allocation between vision and olfaction across the genus *Drosophila*, *Nature Communications* (2019). [DOI: 10.1038/s41467-019-09087-z](https://doi.org/10.1038/s41467-019-09087-z)

Provided by Max Planck Society

Citation: Trade-offs in adaptive anatomy: Vinegar fly species have good vision or olfaction, but not both (2019, March 14) retrieved 9 April 2024 from <https://phys.org/news/2019-03-trade-offs-anatomy-vinegar-species-good.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--