

A warning for wine-lovers: Climate change is messing with your favourite tipple's timing

March 14 2019, by Christopher Davies



Credit: AI-generated image (disclaimer)

While the much-derided "latte set" are stereotyped as the biggest worriers about climate change, it's the chardonnay crowd who are acutely feeling its effects.

Australia's wine industry is both world-renowned and economically



significant, with around A\$5.6 billion in sales in 2016–17, and winemaking and associated tourism responsible for more than 170,000 full and part-time jobs. Statistics also show that wine consumption is now accepted as being just as dinky-di as beer drinking for the average Australian.

However, record-breaking daily maximum temperatures, warmer than average overnight temperatures, and increasingly <u>erratic weather patterns</u> are playing havoc with the way wine grapes grow and ripen. This has knock-on effects for Australian <u>grape</u> growers, wine producers and consumers.

Climate in the vineyard hits the cellar and the store shelf

Most of Australia's wine regions have experienced <u>rising average daily</u> <u>temperatures</u>. One effect is changes to ripening times, which has compressed the harvesting season and given wine-makers a crucial logistical headache.

Traditionally, white grape varieties would generally reach optimum ripeness before red ones. While all grapes tend to ripen faster as temperatures rise, this effect is more pronounced for later-ripening varieties (for example Shiraz and Cabernet Sauvignon) than earlier ripening varieties (for example Chardonnay and Riesling).

The old process of staggered harvesting times for red and white grape varieties was efficient, allowing the winery's capacity to be used in sequence for different varieties. Now that different varieties are ripening at the same time, vineyards and wineries will have to make tough choices about which grapes to prioritise, and which ones to leave until later, resulting in inferior wine. Alternatively, they could take the expensive



decision to increase production capacity by investing in more infrastructure such as fermenters and stainless steel tanks.

Perhaps you're thinking that you, the savvy wine drinker, are unaffected by the difficulties faced by winemakers in the vineyards and wineries far away. Unfortunately this isn't so. Harvesting grapes when they are not at optimal ripeness to solve the logistical problems of processing can lead to lower-value wine.

The fact is that this new reality is costing everyone – grape-growers, winemakers and consumers alike.



Credit: AI-generated image (disclaimer)

And just in case you think that the simple answer is changing Australia's



cultured palates back to beer, think again. Hop production is being <u>hit</u> <u>just as hard by climate change</u>.

Help is at hand

Fortunately, these are problems we hope to tackle. CSIRO recently announced a five-year research partnership with <u>Wine Australia</u>, and one of the projects aims to adjust wine grape ripening to suit a changing climate.

We hope to do it by studying <u>plant growth regulators</u> (PGRs) – molecules that are used by the plant to control and coordinate development. We are using a class of PGRs called auxins, first studied in grass seedlings by Charles Darwin in the 1880s, that have important roles in vine growth, and the timing of grape growth and ripening.

By spraying these compounds onto vines and grapes shortly before ripening, auxins can potentially be used to influence the timing of this process and therefore harvest date. They are already used in other horticultural crops, such as to control fruit drop in apples and pears.

Applying very small amounts of auxin can delay grape ripening, and therefore harvest timing, by up to four weeks (Davies et al., 2015, J Ag Food Chem 63: 2137-2144). This treatment works for red and white varieties in hot or cool climates, and is safe, cheap and easy to apply.

The flavour and aroma of wines made from ripening-delayed grapes is largely indistinguishable from wines made from untreated fruit harvested at the same sugar level, up to a month earlier. An exciting exception is that, in Shiraz, auxin-induced ripening delay can be used to increase the concentration of rotundone, the compound responsible for this variety's popular peppery notes.



Work is currently under way to fine-tune spray formulations and application times. The aim is to release a commercially available product within the next five years.

This kind of solution will be vital for the sustainable, economical production of high-quality wines from existing grape varieties in established wine growing regions. We hope it will ensure you can enjoy your favourite drop for many years to come.

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: A warning for wine-lovers: Climate change is messing with your favourite tipple's timing (2019, March 14) retrieved 17 April 2024 from https://phys.org/news/2019-03-wine-lovers-climate-messing-favourite-tipple.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.