


Bacteria reveal strong individuality when navigating a maze

April 24 2019, by Ori Schipper



Behavioral experiment with bacteria: a T-maze with a chemical gradient presents bacterial cells with the choice of approaching or avoiding the attractant at each branching. Credit: ETH Zurich

Researchers from ETH Zurich demonstrate that genetically identical cells exhibit differing responses in their motility towards chemical attractants. Average values hide the full picture when it comes to describing the behavior of bacteria.

Although they are considered the simplest of all [life forms](#), even microorganisms sense their environment and are able to actively move within it. This allows them to identify both food and harmful substances and to move towards or away from them, guided by the concentration gradient of the substance in their environment. The journey of many microbes can thus be viewed as a sequence of decisions based on chemical gradients.

The ability of cells to target or avoid particular substances is called [chemotaxis](#). Until now, scientists have generally considered the chemotactic properties of [bacteria](#) to be a common feature of a species or population—as if all cells behaved more or less the same. In this case, average values are sufficient to describe their movement behaviour. Now, researchers at ETH Zurich have observed the chemotaxis of bacteria in a behavioural experiment. "If you look with the appropriate technology, you'll find astonishing behavioural differences even within a population of genetically identical cells," report Mehdi Salek and Francesco Carrara, the lead authors of a study recently published in *Nature Communications*.

Microbes in the T-maze

Together with their colleagues in the research group led by Professor Roman Stocker at the Institute of Environmental Engineering, they have developed a special microfluidic system that allows them to observe the movement of thousands of individual bacteria in a liquid at extremely small scales. The system comprises a series of narrow channels that branch out onto a thin glass plate to form a sort of microscopic maze through which the bacteria swim.

Such mazes are often used in experimental studies of the behavioural preferences of other organisms, such as insects or worms (and also plant roots). With their microfluidic system, the ETH researchers were able

for the first time to apply this traditional tool of ecologists on a microscopic scale. Their maze resembles a family tree, with a starting channel that branches out again and again towards the bottom, where the concentration of a chemical attractant is at its highest.

Decisions at the fork in the road

The bacteria all start in the same place—and visibly divide up within the channel system as they are forced to decide at each fork whether to swim up or down the gradient of attractant. The bacteria owe their chemotactic abilities to specialised receptors that allow them to identify the attractant. In addition, they have about half a dozen flagella, which can rotate either clockwise or anti-clockwise. "Based on this, the bacterium changes its direction or continues to swim in one direction," explain Salek and Carrara.

Even within a group of genetically [identical cells](#)—that is, clones—the ETH researchers found individuals that were able to follow the attractant well (by navigating towards the higher concentration whenever they came to a fork), and those that were less able to negotiate the maze. The scientists attribute these behavioural differences to variations in the genetic activity of identical genes in sister cells. This means the cells have different amounts of the corresponding proteins. "There is biochemical noise in every cell. As a fundamental random component, this causes diversity of appearance and behaviour," say the researchers.

Successful population of individualists

Diversity, or heterogeneity, of chemotaxis may provide an evolutionary advantage for the bacteria, since although those skilled at chemotaxis can quickly locate and exploit locally stable food sources, their sister [cells](#) less affected by the attractant are more likely to venture into new

territory, where they may encounter additional food sources in a constantly changing environment.

"Non-genetic diversity has long been known in the biomedical life sciences; for example, it is thought to play a role in antibiotic resistance. Now, environmental scientists have shown that this [diversity](#) also affects fundamental behaviours of bacteria, such as locomotion and chemotaxis—further expanding the concept of bacterial individuality," says Stocker. He assumes that the varying individual behaviour of bacteria may also be relevant to gaining a better understanding of processes such as the pathogenic infection of corals or the bioremediation of oil spills.

More information: M. Mehdi Salek et al, Bacterial chemotaxis in a microfluidic T-maze reveals strong phenotypic heterogeneity in chemotactic sensitivity, *Nature Communications* (2019). [DOI: 10.1038/s41467-019-09521-2](#)

Provided by ETH Zurich

Citation: Bacteria reveal strong individuality when navigating a maze (2019, April 24) retrieved 27 April 2024 from

<https://phys.org/news/2019-04-bacteria-reveal-strong-individuality-maze.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.