

Bacteria uses viral weapon against other bacteria

April 25 2019



The virus SW1 is carried by cells so that related bacteria may recognize other SW1 carriers and kill bacteria that do not have the virus, giving bacteria with SW1 a competitive advantage when foraging for food. Credit: Sooyeon Song and Missy Hazen/Penn State



Bacterial cells use both a virus—traditionally thought to be an enemy—and a prehistoric viral protein to kill other bacteria that competes with it for food according to an international team of researchers who believe this has potential implications for future infectious disease treatment.

The team, led by Thomas Wood, holder of the biotechnology endowed chair and professor of chemical engineering, Penn State, noticed a gap between two kinds of bacterial cells as they moved on agar plates toward each other, in a form of motility known as "swimming" where the bacterium moves by rotating its flagella.

"It was kind of like when Alexander Fleming noticing <u>bacteria</u> did not grow near a fungus when he discovered penicillin in the 1920s," Wood said. "We then investigated and found the gap was caused by cells being killed by a virus, SW1, that was carried by only one of the strains."

The team not only found that a bacterial cell uses the SW1 virus to kill bacteria that are its competitors for food, they also discovered that bacteria that carry the virus are more fit and not killed as often as bacteria without the virus. In addition, they found that for a bacterium to be able to use SW1, it needs a protein fossil of a bacterium that came from another virus millions of years ago. They report their results in a recent issue of *Cell Reports*.

Wood explained that sometimes a virus will enter a cell and hide in the chromosome until it is ready to attack and kill the cell. However, due to random mutations, a virus sometimes is stuck in the chromosome and cannot leave to attack. There are nine viruses trapped in this way in the beneficial gut bacterium E. coli. To be activated and used by E. coli against other bacteria, the virus SW1 needs a protein, YfdM, from a virus that became caught in the E. coli chromosome millions of years ago.



In addition, the virus is used by the bacterium as an identifier.

"Bacteria are frequently thought of as living alone but instead they can forage for food as groups. In order to act as a group, they must be able to distinguish themselves from other bacteria," Wood said. "In one type of social activity, <u>bacterial cells</u> secrete chemical signals to communicate. But now, we show that bacterium cells use viruses to distinguish themselves from closely-related bacteria."

If the bacterium does not detect the virus in the other <u>bacterium</u>, it identifies a competitor for food, and unleashes the <u>virus</u> to kill its competition. While the exact workings of this process are not yet understood, it may hold <u>potential benefits</u> for research and medicine.

"Understanding how cells compete is important," Wood said. "For example, in <u>synthetic biology</u> where many cells work together, you can make just the <u>cells</u> you want to work for you become dominant. In medicine, you can devise better ways to fight infectious agents."

Provided by Pennsylvania State University

Citation: Bacteria uses viral weapon against other bacteria (2019, April 25) retrieved 11 May 2024 from <u>https://phys.org/news/2019-04-bacteria-viral-weapon.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.