


Carbon dioxide to methanol conversion

April 5 2019



(Left) Schematic showing the production of methanol by electrocatalytic reduction of carbon dioxide using the PD-Zn/Ag foam catalyst. (Right) Scanning electron microscopy image of the surface of the PD-Zn/Ag foam catalyst. The areas with “holes” are the silver foam while the other parts show the zinc dendrites lying on it. Credit: National University of Singapore

NUS chemists have developed a highly efficient nanostructured catalyst based on zinc and silver that can convert carbon dioxide, an

environmental pollutant and greenhouse gas, to methanol for use as chemical feedstock and fuel.

The electrocatalytic reduction of [carbon dioxide](#) using [renewable electricity](#) and an appropriate [catalyst](#) is a promising green manufacturing method to produce chemicals and fuels sustainably. Methanol is one of the most valuable products that can be generated from this process. Apart from its use as a fuel, it is also employed as a chemical building block to produce more complex chemicals such as acetic acid. While the chemical structure for methanol may be simple, its conversion efficiency from carbon dioxide is poor.

A research team led by Prof Yeo Boon Siang, Jason, from the Department of Chemistry, NUS in collaboration with Dr. Federico Calle-Vallejo from the University of Barcelona, Spain has discovered that zinc dendrites deposited on silver foam (termed as PD-Zn/Ag foam), can be used as a catalyst to convert carbon dioxide to methanol with high efficiency. Zinc and silver metals by themselves, and their alloys, are more effective for converting carbon dioxide to carbon monoxide. However, by patterning them at the nanometer scale, their functionality as a catalyst can be enhanced. The resulting PD-Zn/Ag foam is able to produce methanol with a Faradaic efficiency and current density reaching as high as 10.5% and -2.7mA/cm^2 , respectively. This represents a tenfold increase over conventional zinc-silver catalysts. Using [experimental results](#) and theoretical calculations, the catalytically active sites were identified to be strained zinc dendrites deposited on the silver supporting material. These active sites bind to carbon monoxide intermediates strongly, which in turn facilitate their conversion to methanol.

Prof Yeo said, "This work illustrates that nanostructured bimetallic systems can enhance both the activity and selectivity of the catalytic [carbon](#) dioxide reduction reaction. The research outcome can be

exploited to design and synthesise catalysts with improved functionality."

Building on the research findings from their work, the research team plans to develop catalysts with higher methanol conversion efficiency.

More information: Qi Hang Low et al. Enhanced Electroreduction of Carbon Dioxide to Methanol Using Zinc Dendrites Pulse-Deposited on Silver Foam, *Angewandte Chemie International Edition* (2018). [DOI: 10.1002/anie.201810991](https://doi.org/10.1002/anie.201810991)

Provided by National University of Singapore

Citation: Carbon dioxide to methanol conversion (2019, April 5) retrieved 2 May 2024 from <https://phys.org/news/2019-04-carbon-dioxide-methanol-conversion.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--