

The emergence of male and female traits in the development of beetle horns

April 11 2019



Male Japanese rhinoceros beetle (left), female beetle (center), Knockdown phenotypes of transformer gene (right). Male-like horns are formed in the female beetle. Credit: NIBB

The male Japanese rhinoceros beetle, *Trypoxylus dichotomus*, which lives on Japan's main island, has big horns that are used as weapons to fight other males for females. Scientists have sought the developmental mechanism that creates these horns, and to this end, a research team at the National Institute for Basic Biology in Japan has identified sex-determining genes for the rhinoceros beetle, and has succeeded in identifying the timing of sex differences that appear in horn primordia.

A member of the research team, Dr. Shinichi Morita, said, "Unlike the model animal *Drosophila*, we did not have any basic information concerning these [beetles](#), such as their developmental stages. Furthermore, because the larvae and pupae live in the soil, it was particularly difficult to observe their development." Therefore, Dr. Morita developed a rearing method encompassing the beetle's late larval stage up until adulthood that used plastic tubes, but didn't require any soil. Dr. Morita then video recorded a few weeks of the process of beetle larvae becoming pupae, and looked for signs that the prepupal stage had started.

"We found that head-rocking behavior was a sign of the beginning of the prepupal stage," Dr. Morita said. Using this as a reference point, the research group was able to create a stage table of [horn](#) formation along a time axis. As a result of this, it became clear that it took about 36 hours after the reference point for the morphological differences between males and females in the horn primordia to appear.



Late larval stage (left) , prepupal stage (center) and pupal stage (right). Credit: NIBB

Next, referring to the mechanism of sex determination in *Drosophila*, they examined how the sex of the beetle was determined. The research team subsequently revealed that the sex of the beetle is controlled by a gene called Transformer. They found that when this gene is suppressed during the larval stage, male-like horns are formed in the female beetle.

Upon this discovery, the researchers then looked for a boundary between the time when the horn can be grown by the suppression of transformer gene and the time when the horn cannot be grown, by suppressing this gene in female larvae using RNAi taken from various stages during

larval-pupal development. They ultimately succeeded in identifying the time when the sex difference of this horn appeared.

The leader of the research team, Prof. Teruyuki Niimi, said, "The timing at which the gene that causes sexual differences in horn formation is about 29 hours after the baseline point of the prepupal stage start. Based on this finding, we want to clarify the details of the gene network of horn formation."

The results of this were published on April 10, 2019 in *PLOS Genetics*.

More information: "Precise staging of beetle horn formation in *Trypoxylus dichotomus* reveals the pleiotropic roles of doublesex depending on the spatiotemporal developmental contexts" by Shinichi Morita, Toshiya Ando, Akiteru Maeno, Mutsuki Mase, Takeshi Mizutani, Shuji Shigenobu, Teruyuki Niimi were published in *PLOS Genetics*. [DOI: 10.1371/journal.pgen.1008063](https://doi.org/10.1371/journal.pgen.1008063)

Provided by National Institutes of Natural Sciences

Citation: The emergence of male and female traits in the development of beetle horns (2019, April 11) retrieved 17 July 2024 from <https://phys.org/news/2019-04-emergence-male-female-traits-beetle.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.