

Genetic breakthrough on tropical grass could help develop climate-friendly cattle farms

April 9 2019



Study author Margaret Worthington examines *Brachiaria* grasses in an experimental field at the International Center for Tropical Agriculture near Cali, Colombia. Credit: Paul Blake / International Center for Tropical Agriculture

Cattle are a mainstay for many smallholders but their farms are often on



degraded lands, which increases cattle's impact on the environment and lowers their production of milk and meat. Researchers at the International Center for Tropical Agriculture (CIAT) have shown that *Brachiaria* grass species can reduce greenhouse gas emissions from cattle and increase productivity—and breeding improved varieties can potentially augment the environmental and economic benefits.

But the breeding process is difficult, time-consuming and expensive. A breakthrough on *Brachiaria*'s complex genome may make breeding much more efficient, and potentially increase the speed with which new grasses begin benefiting cattle farmers and the environment.

Margaret Worthington, a geneticist at CIAT and the University of Arkansas, and colleagues created the first dense molecular map of *B. humidicola*, a robust and environmentally friendly forage grass. They also pinpointed the candidate genes for the plant's asexual reproductive mechanism, which is a huge asset for plant breeders. The findings were published in January in *BMC Genomics*.

"The idea is to create a better crop with less time and less money and to get it out faster to farmers," said Worthington. "By using this molecular marker, you increase the odds of finding that rare winner."

Traditional plant-breeding methods for *Brachiaria* grasses involve one of two complex techniques. One is to grow the plant to seed, and to study the seeds under a microscope to determine if the plant reproduced asexually. The other involves excising the plant's embryos and conducting a similar analysis. Both techniques require many weeks, significant funds and highly trained specialists.

Asexual reproduction through seed, called apomixis, is key for developing new crop varieties for widespread use. Crops that reproduce through apomixis conserve the same traits from one generation to the



next, essentially locking in sought-after characteristics such as drought tolerance or high nutritional value. Plants that reproduce sexually do not reliably pass on desired traits to subsequent generations.



Study author Margaret Worthington examines *Brachiaria* grasses in an experimental field at the International Center for Tropical Agriculture near Cali, Colombia. Credit: Paul Blake / International Center for Tropical Agriculture

Seeds, perpetually

With this molecular marker, plant breeders can run a quick and inexpensive test when *Brachiaria* grasses are seedlings to identify whether they reproduce through apomixis. The results are available in a



couple of weeks. This allows plant breeders to select only asexually reproductive plants for trials, allowing them to allocate more time and resources to <u>plants</u> that have the potential to produce new cultivars.

Brachiaria grasses have often been considered an "orphan crop," due to a lack of investment in research, but their potential for making tropical farms more productive and better for the environment is well known among tropical forage specialists. One recent study found that *B. humidicola* was especially adept at reducing the <u>nitrous oxide</u>, a strong greenhouse gas, emitted from soil as result of cattle urine deposition. In addition, CIAT researchers have identified mechanisms that this tropical grass uses to efficiently acquire nutrients from soil.

Brachiaria breeders also value apomixis for smallholders in developing nations who have limited resources for investing in improving their farms. Improved grass varieties that produce sufficient quantities of traitretaining seeds can eliminate the need to purchase new seeds for every planting, which is a potentially expensive barrier to adoption.

"This breakthrough allows for the acceleration of our breeding program for multiple traits, including the development of tropical forages that can help reduce <u>greenhouse gas emissions</u> and make farming more ecoefficient," said Joe Tohme, a senior scientist at CIAT and study coauthor.

"This discovery represents a milestone in the path toward developing mitigation technologies in the livestock production sector," said Jacobo Arango, a study co-author who is an environmental biologist from CIAT and a Lead Author for the next Assessment Report on Climate Change Mitigation of the Intergovernmental Panel on Climate Change (IPCC).

More information: Margaret Worthington et al, Translocation of a parthenogenesis gene candidate to an alternate carrier chromosome in



apomictic Brachiaria humidicola, *BMC Genomics* (2019). <u>DOI:</u> <u>10.1186/s12864-018-5392-4</u>

Provided by International Center for Tropical Agriculture (CIAT)

Citation: Genetic breakthrough on tropical grass could help develop climate-friendly cattle farms (2019, April 9) retrieved 19 April 2024 from <u>https://phys.org/news/2019-04-genetic-breakthrough-tropical-grass-climate-friendly.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.