

Melting glaciers causing sea levels to rise at ever greater rates

April 8 2019



Melting glaciers in Russia's Arctic territory. This image shows the blue glaciers on the reddish-brown Franz Josef Land archipelago. The glaciers (blue) are covered with little or no snow (white), which is a clear indicator of significant loss of mass. Credit: Copernicus Sentinel data 2017

Melting ice sheets in Greenland and the Antarctic as well as ice melt from glaciers all over the world are causing sea levels to rise. Glaciers alone have lost more than 9 trillion tons of ice since 1961, raising water



levels by 27 millimeters, an international research team under the lead of the University of Zurich has now found.

Glaciers have lost more than 9 trillion tons (that is 9,625,000,000,000 tons) of ice between 1961 and 2016, which has resulted in global sea levels rising by 27 millimeters in this period. The largest contributors were glaciers in Alaska, followed by the melting ice fields in Patagonia and glaciers in the Arctic regions. Glaciers in the European Alps, the Caucasus and New Zealand were also subject to significant ice loss; however, due to their relatively small glacierized areas, they played only a minor role when it comes to the rising global sea levels.

Combination of field observations and satellite measurements

For the new study, the international research team combined glaciological field observations with geodetic <u>satellite measurements</u>. The latter digitally measure the surface of the Earth, providing data on ice thickness changes at different points in time. The researchers were thus able to reconstruct changes in the ice thickness of more than 19,000 glaciers worldwide.

The researchers used a comprehensive database compiled by the World Glacier Monitoring Service maintained by a worldwide network of observers. The researchers combined this data with their own satellite analyses. "By combining these two measurement methods and having the new comprehensive dataset, we can estimate how much ice has been lost each year in all mountain regions since the 1960s," explains Michael Zemp, who led the study. "The glaciological measurements made in the field provide the annual fluctuations, while the satellite data allows us to determine overall ice loss over several years or decades."





Regional share of glaciers in sea-level rise from 1961 to 2016. Credit: Zemp

335 billion tons of ice lost each year

The global mass loss of glacier ice has increased significantly in the last 30 years, and currently amounts to 335 billion tons of lost ice each year. This corresponds to an increase in sea levels of almost one millimeter per year. "Globally, we lose about three times the ice volume stored in the entirety of the European Alps—every single year," says glaciologist Zemp. The melted ice of glaciers therefore accounts for 25 to 30 percent of the current increase in global sea levels. This ice loss of all glaciers roughly corresponds to the mass loss of Greenland's <u>ice sheet</u>, and



clearly exceeds that of the Antarctic.

More information: Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016, *Nature* DOI: 10.1038/s41586-019-1071-0, www.nature.com/articles/s41586-019-1071-0

Provided by University of Zurich

Citation: Melting glaciers causing sea levels to rise at ever greater rates (2019, April 8) retrieved 10 May 2024 from <u>https://phys.org/news/2019-04-glaciers-sea-greater.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.