

An important function of non-nucleated sperm

April 29 2019



Eupyrene sperm (upper photo) and an apyrene sperm (lower photo) of of the silk moth, *Bombyx mori*. Credit: NIBB

Some animals form characteristic infertile spermatozoa called parasperm, which differ in size and shape compared to fertile sperm produced by single males. Species that have been reported to produce parasperm include snails, cottoid fish, moths and butterflies. Moths and butterflies produce fertile eupyrene sperm and anucleate non-fertile parasperm, which are known as apyrene sperm. A research team at the National Institute for Basic Biology in Japan has identified the gene

involved in the formation of the apyrene sperm and has revealed the important function of the apyrene sperm in fertilization using the silk moth, *Bombyx mori*.

A member of the research team, Dr. Hiroki Sakai said, "Parasperm has been discovered in multiple animals despite its inability to be fertilized itself. From this, there is a possibility that parasperm may play an important role in fertilization." Although many morphological observations and ecological surveys have been conducted on parasperm, genes involved in the formation of parasperm have not been identified in any species.

The researchers found that Sex-lethal (Sxl), which is a widely studied key sex determination gene in the fruit fly *Drosophila melanogaster*, is an essential gene for the formation of apyrene [sperm](#) in the silk [moth](#). "The Sxl gene plays an important role in female determination in the fruit fly. We are surprised that the gene was involved in the formation of non-nucleated sperm," Dr. Sakai said.

The research group has created a silk moth whose Sxl gene function has been inhibited by the genome editing technology. As a result, they demonstrated that this [silk](#) moth could not form accurate apyrene sperm. Furthermore, the mating experiment using Sericology techniques revealed that its eupyrene sperm is normal. Based on this, it was revealed that Sxl is required for the formation of apyrene sperm.

Silk moth males, whose Sxl gene does not work, cannot produce future generations because they cannot produce apyrene sperm. Why can't they reproduce without apyrene sperm? The research team has clearly shown that apyrene sperm is required for the eupyrene sperm migration in the female organs.

The leader of the research team, Prof. Niimi said, "The significance of

this study is that it is the first time that specific [genes](#) required for parasperm formation have been identified. Moreover, we demonstrated that parasperm is necessary for eupyrene sperm migration in female organs. The results indicate that anucleate parasperm, while appearing less important in fertilization actually are so, may have a major impact on future sperm research in animals."

The results of the study will be published on the week of April 29, 2019 in *Proceedings of the National Academy of Sciences*.

More information: Hiroki Sakai et al., "Dimorphic sperm formation by Sex-lethal," *PNAS* (2019).

www.pnas.org/cgi/doi/10.1073/pnas.1820101116

Provided by National Institutes of Natural Sciences

Citation: An important function of non-nucleated sperm (2019, April 29) retrieved 18 April 2024 from <https://phys.org/news/2019-04-important-function-non-nucleated-sperm.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--