

Quality of laser beam shaping can be enhanced at no extra cost

April 24 2019, by Osaka University




Figure 1. Square flattop beam profiles made by: (a) Diffractive Optical Element (DOE), (b) vertical phase grating (conventional method), (c) virtual diagonal phase grating (new method). Credit: © 2019 Nakata Y. et al., *Scientific Reports*.

Researchers from Osaka University have developed a technique for improving accuracy of laser beam shaping and wavefront obtained by conventional methods with no additional cost by optimizing virtual phase grating. The results of their research were published in *Scientific Reports*.

A high quality square flattop [beam](#) is in demand for various fields, such as uniform laser processing and medicine, as well as ultrahigh intensity laser applications for accelerators and nuclear fusion. Beam [shape](#) is key to realizing the laser's potential abilities and effects. However, since beam shape and wavefront vary by laser, beam shaping is essential for producing the desired shapes to respond to various needs.

Static and adaptive beam shaping methods have been developed for various applications. With Diffractive Optical Element (DOE) as a static method, edge steepness and flatness are low and wavefront becomes deformed after shaping. (Figure 1 (a)) In addition, computer-generated hologram (CGH) as a typical adaptive method has the same difficulties.

Meanwhile, an adaptive beam shaping technique that uses [phase](#) grating encoded on a [spatial light modulator](#) (SLM) with spatial-frequency filtering in the Fourier plane in a 4f system was developed. (Figure 2 (a)) This conventional method generates a square flattop beam by spatially controlling diffraction efficiency without deforming the wavefront. However, because the extracted and residual components overlap in the Fourier plane, it was necessary to cut the high spatial-frequency (HSF) [component](#) from the extracted component, limiting the flatness and the edge steepness of the resultant beam shape. (Figure 1 (b))




Figure 2. Experimental layout: phase grating and filtering in the Fourier plane of the 4f system. (a) vertical phase grating (conventional method), (b) virtual diagonal phase grating (new method). Credit: © 2019 Nakata Y. et al., *Scientific Reports*.

In this study, the group developed a universal beam shaping technique at [high accuracy](#), which can be used for various lasers from ultraviolet to near-infrared domain.

This method spatially separates the residual and extracted components in the Fourier plane by using a virtual diagonal phase grating (Figure 2(b)) and clears overlap by making the grating vector, k_g , non-parallel to the normal vectors, k_x or k_y , of the desired beam profile, which are parallel to each other in the conventional scheme.

By efficiently using only extracted components containing HSF

components, beam shaping at high resolution was achieved. This allowed for a highly uniform flattop beam of any cornered shape without ripples, suppressing the edge of the shaped beam to a height of 20 μm , which is less than 20% of that obtained with conventional vertical phase grating.

Corresponding author Yoshiki Nakata says, "Our [method](#), which allows for optimization of beam shaping by improving resolution and accuracy, will contribute to a wide field, including basic research, manufacturing and medical engineering. In conventional beam shaping systems, beam shaping accuracy can be significantly enhanced at no extra cost simply by changing the spatial frequency filter and phase [grating](#) encoded on an SLM."

More information: Yoshiki Nakata et al. Utilization of the high spatial-frequency component in adaptive beam shaping by using a virtual diagonal phase grating, *Scientific Reports* (2019). [DOI: 10.1038/s41598-019-40829-7](#)

Provided by Osaka University

Citation: Quality of laser beam shaping can be enhanced at no extra cost (2019, April 24) retrieved 27 April 2024 from <https://phys.org/news/2019-04-quality-laser-extra.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--