


Rusted root: Weedy rice repeatedly evolves 'cheater' root traits

April 5 2019, by Talia Ogliore



New imaging techniques allow researchers to consider how weedy rice plants have repeatedly evolved root systems that may convey a competitive advantage. Credit: Christopher Topp, Donald Danforth Plant Science Center

Weedy rice is neither wild rice nor crop rice, but rice gone rogue that

has shed some traits important to people. It also is an incredibly aggressive, potentially detrimental weed that pops up almost everywhere rice is grown, and it can reduce crop yields by more than 80 percent if it invades a field.

Researchers from Washington University in St. Louis and the Donald Danforth Plant Science Center used a new imaging technique to reveal a takeover strategy that has worked for weedy rice over and over again: roots that minimize below-ground contact with other plants.

"Weedy rice may have evolved a go-it-alone 'cheater' root growth strategy that could allow it to exploit the nutrient-sharing soil environment of rice fields," said Kenneth M. Olsen, professor of biology in Arts & Sciences at Washington University and senior author on a new paper in *New Phytologist* relating their findings.

"We tend to think of competition occurring above ground because that's the part of the plant we see. But that's only half the plant," Olsen said. "It's the 'hidden half'—i.e., the root system—that plays a critical role in some of the most important aspects of plant growth and survival, including water uptake and competition for [essential nutrients](#) like nitrogen and phosphorous."

By some estimates, root growth is actually a more important determinant of competitive success than above-ground growth, Olsen said.

"This appears to be particularly true for agricultural weeds such as weedy rice, which compete against crop varieties in agricultural fields," he said.

A look underground

Scientists are only just beginning to get their first realistic glimpses of

how root growth and below-ground root interactions affect a plant's ability to compete for resources. In the past, they couldn't get a decent look at a root system without digging it up, which inevitably damages it, or by growing plants in very artificial conditions, such as sandwiched between two glass plates.

For this study, researchers compared the roots of two independently evolved types of weedy rice that occur in the same rice fields in the southern United States. Using new imaging techniques, including a semi-automated optical tomography approach developed by Christopher Topp at the Donald Danforth Plant Science Center, the researchers took more than 70 photographs of the root systems of each of 671 different weedy rice [plants](#). Then they modeled the pictures in 3-D to create 360-degree digital maps of their roots.

The researchers used a variety of algorithms they developed to analyze 98 [physical traits](#)—including root depth, [root system](#) width, certain exploratory traits and root-soil angles.

They also conducted [genetic analyses](#) to track the weeds' separate paths from their domesticated pasts to their persona-non-grata status in the rice fields of today.

"Natural selection says that they (the two types of weedy rice) should respond to this environment by evolving similar traits," said Marshall J. Wedger, a Ph.D. candidate in Olsen's laboratory and first author on the paper. "They did evolve similar traits in response to similar environmental pressures, but they did so using very different genetic mechanisms."

The new study shows how two independently evolved weedy rice strains have convergently arrived at a similar pattern of [root](#) growth that may play a role in their ability to outcompete cultivated [rice](#) for soil nutrients.

"By looking at the genetic basis of [weedy rice](#) evolution, we can see whether—at the genetic level—there's more than one way to evolve a weed," Olsen said. "What we find, both for aboveground traits and now with this study for belowground traits, is that the answer is a definitive yes."

"In other words, it's disconcertingly easy to evolve a weed from a domesticated crop," Olsen said. "This can occur multiple times independently from different crop varieties."

More information: Marshall J. Wedger et al, Convergent evolution of root system architecture in two independently evolved lineages of weedy rice, *New Phytologist* (2019). [DOI: 10.1111/nph.15791](https://doi.org/10.1111/nph.15791)

Provided by Washington University in St. Louis

Citation: Rusted root: Weedy rice repeatedly evolves 'cheater' root traits (2019, April 5) retrieved 5 May 2024 from <https://phys.org/news/2019-04-rusted-root-weedy-rice-repeatedly.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--