


Tracking charge carriers in the molecular crystal at organic pn junction

April 26 2019



(a) Hopping conduction in disordered junction, (b) band conduction in crystalline junction. Credit: NINS/IMS

In conventional organic solar cells, the electrons exhibit their particle-nature and need to jump between organic molecules in the cell. The conductivity is, therefore, lower than that of crystalline silicon solar cells. Researchers have succeeded in arranging the organic molecules in a highly ordered manner like in crystals, and to invoke the wave-nature. "Conductive bands" are formed by energy dispersive states and contribute to the high-carrier conductivity. It may improve the total efficiency of the cell.

An organic solar cell is a light, flexible, low cost and green device, hence regarded as a potential seed of innovation in the renewable energy industry. The efficiency of energy conversion of the organic solar cell is, however, lower than those of contemporary silicon [solar cells](#).

In semiconductor solar cells, the light is converted into an energized pair of an electron (negative carrier) and a hole (positive carrier) at the "pn junction" interface at two semiconductor layers in the cell. Donor (electron pitching; [p-type](#)) and acceptor (electron catching; n-type) molecules in each layer of the semiconductors face each other as the ideal p/n junction. To increase the number of such solar "batteries" in the cell, a large area of pn junction is required, so that a complicated "bulkhetero" pn junction, which is a folded interface like pleats, has been developed. In such a complicated structure like a maze, the generated carriers are difficult to reach output electrodes of the cell because the molecules are arranged roughly, in other words, crystallinity is low (Fig. 1(a)). To realize a high transport efficiently, the carrier, an electron or a hole, should delocalize between molecules as a matter wave (Fig. 1(b)). The ordered arrangement of molecules brings out the wave-nature of the carries.

Researchers at the Institute for Molecular Science (IMS), Japan Synchrotron Radiation Research Institute (JASRI) and Tokyo University of Science have succeeded in fabricating the organic semiconductor pn junction with high crystallinity. In the fabrication process of the junction, the acceptor (perfluoropentacene) molecules were deposited in a well-ordered manner onto the single crystal of the donor molecules (pentacene) by the [molecular beam epitaxy](#) (MBE) technique. The electronic structures of the crystalline pn junction were observed by angle-resolved [photoelectron spectroscopy](#) and showed the layer of the acceptor molecules form the valence band which is evidence of invoking the wave-nature. The result of the present study shows that the MBE facilitates the fabrication of the crystalline pn junction which may bring

out the wave nature of both the electrons and the holes.

The functions of organic semiconductors can be tuned by designing the structures of constituent organic molecules. The fabrication technology of the crystalline pn junctions using a variety of [organic molecules](#) enables us to realize the new concept organic solar [cells](#) with high energy conversion efficiency.

More information: Yasuo Nakayama et al, Widely Dispersed Intermolecular Valence Bands of Epitaxially Grown Perfluoropentacene on Pentacene Single Crystals, *The Journal of Physical Chemistry Letters* (2019). [DOI: 10.1021/acs.jpclett.8b03866](https://doi.org/10.1021/acs.jpclett.8b03866)

Provided by National Institutes of Natural Sciences

Citation: Tracking charge carriers in the molecular crystal at organic pn junction (2019, April 26)
retrieved 3 May 2024 from

<https://phys.org/news/2019-04-tracking-carriers-molecular-crystal-pn.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.