

New research supports volcanic origin of Kiruna-type iron ores

April 12 2019



Credit: CC0 Public Domain

The origin of so-called Kiruna-type apatite-iron oxide ores has been the topic of a longstanding debate for over 100 years. In a new article published in *Nature Communications*, a team of scientists presents new

and unambiguous data in favour of a magmatic origin for these important iron ores. The study was led by researchers from Uppsala University in Sweden.

Despite globally increasing demand for rare metals, iron is the overall most important metal for modern industry. Over 90 percent of Europe's total iron production comes from apatite-iron oxide ores, also referred to as Kiruna-type ores, named after the extremely large and iconic Kiruna iron ore deposit in northern Sweden. Today, the Swedish deposits at Kiruna and Malmberget are the largest and most important iron producers in Europe, and Kiruna-type deposits represent an iron source of global importance. These deposits also have a large future potential for production of sought-after and critical rare earth elements (REE) as well as phosphorous, another element deemed critical for Europe's future development.

The [origin](#) and actual process of formation of Kiruna-type ores has remained highly controversial for over 100 years, with suggestions including a purely low-temperature hydrothermal origin, sea floor precipitation, a high-temperature volcanic origin from magma, and high-temperature magmatic fluids. To clarify the origins of Kiruna-type ores, a team of scientists from Uppsala University, the Geological Survey of Sweden, the Geological Survey of Iran, the Indian Institute of Technology in Bombay, and the Universities of Cardiff and Cape Town, led by Uppsala researcher Prof. Valentin Troll, employed Fe and O isotopes, the main elements in magnetite (Fe_3O_4), from Sweden, Chile and Iran to chemically fingerprint the processes that led to formation of these ores.

By comparing their data from Kiruna-type iron ores with an extensive set of magnetite samples from [volcanic rocks](#) as well as from known low-temperature hydrothermal iron ore deposits, the researchers were able to show that more than 80 percent of their magnetite samples from Kiruna-

type apatite-iron oxide ores were formed by high-temperature magmatic processes in what must represent volcanic to shallow sub-volcanic settings. The new results constitute an important advance in our understanding of Kiruna-type ores and will be of help for the interpretation of, and future exploration for, apatite-[iron](#) oxide deposits worldwide.

More information: V. R. Troll et al. 2019: Global Fe-O isotope correlation reveals magmatic origin of "Kiruna-type" apatite-iron oxide ores. *Nature Communications*, [DOI: 10.1038/s41467-019-09244-4](https://doi.org/10.1038/s41467-019-09244-4)

Provided by Uppsala University

Citation: New research supports volcanic origin of Kiruna-type iron ores (2019, April 12) retrieved 3 May 2024 from <https://phys.org/news/2019-04-volcanic-kiruna-type-iron-ores.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
