

ATLAS Experiment adds more pieces to the Higgs boson puzzle

May 7 2019





Cross sections time branching fraction for the main Higgs production modes at the LHC (ggF, VBF, VH and ttH+tH) in each relevant decay mode ($\gamma\gamma$, WW, ZZ, $\tau\tau$, bb). All values are normalized to Standard Model predictions. In addition, the combined results for each production cross-section are also shown, assuming the Standard Model values for the branching ratios into each decay



mode. Credit: ATLAS Collaboration/CERN

The Higgs boson was discovered in 2012 by the <u>ATLAS</u> and <u>CMS</u> Experiments at CERN, but its coupling to other particles remains a puzzle.

Fortunately, the LHC provides many windows into measuring Higgs boson couplings. There are four main ways to produce the Higgs boson: through the fusion of two gluon particles (gluon-fusion, or ggF), through the fusion of weak vector bosons (VBF), or in association with a W or Z boson (VH), or one or more top quarks (ttH+tH). There are also five main channels in which Higgs bosons can decay: into pairs of photons, W or Z bosons, tau leptons or b quarks. Each of these processes brings unique insights into the Higgs boson properties.

Thanks to the unprecedented quantities of Higgs bosons produced at the LHC, all of the above production and decay modes have now been observed. In a <u>new result presented by the ATLAS Collaboration</u>, using data collected through 2017, the measurements for each of these processes have reached the significance threshold of five standard deviations, beyond which their existence is considered established.

The Higgs boson yields for most of the combinations of production and decay have been measured (see figure) and have been found to agree with Standard Model predictions. The measurement of the cross sections for each production mode in proton–proton collisions at 13 TeV, assuming the decays occur as predicted by the Standard Model, are the most precise ones obtained to date.

Physicists have also begun to explore the Higgs boson puzzle in a new way. In the latest analyses, instead of counting Higgs bosons inclusively



in the major production and decay modes, ATLAS physicists have measured Higgs boson topologies separately for smaller regions of phasespace: different ranges of Higgs boson transverse momentum, numbers of associated jets, and numbers and kinematic properties of associated weak bosons and top quarks. Using these smaller puzzle pieces, called "simplified template cross sections" (STXS), allows physicists to better separate the measurement process from the interpretation in terms of theoretical properties. Ultimately, it provides a finer-grained picture of Higgs boson couplings at the LHC and more stringent tests of the Standard Model.

Among the STXS regions considered in the analysis, some have already been measured with good precision at the LHC, but no deviation from the Standard Model has been observed so far. These measurements allow physicists to further enhance the sensitivity on the coupling properties of the Higgs boson to the other elementary particles. Further, they have set constraints on new physics theories – such as the "two-Higgs doublet <u>model</u>", which introduces additional Higgs bosons, and the hMSSM supersymmetric model – which are more stringent than those reported previously by ATLAS.

These measurements will continue to improve as more data from Run 2 and beyond are included, providing a yet-finer picture of the properties of the Higgs <u>boson</u>.

More information: Combined measurements of Higgs boson production and decay using up to 80 fb–1 of proton—proton collision data at 13 TeV collected with the ATLAS experiment: atlas.web.cern.ch/Atlas/GROUPS ... ATLAS-CONF-2019-005/

Provided by ATLAS Experiment



Citation: ATLAS Experiment adds more pieces to the Higgs boson puzzle (2019, May 7) retrieved 2 May 2024 from <u>https://phys.org/news/2019-05-atlas-pieces-higgs-boson-puzzle.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.