

Manipulating atoms one at a time with an electron beam

May 17 2019, by David L. Chandler



Illustration of competing experimental P dopant dynamics in graphene and its control. The frames are medium-angle annular dark-field images, and the chemical identity of each dopant was confirmed by electron energy-loss spectroscopy (EELS). (A) Three frames showing a direct exchange between the brighter (due to its greater scattering contrast) P atom and a C neighbor, with the initial (frame 1), transition (frame 2), and final configurations (frame 3). White and black dashed lines indicate the row of the scanning beam when the exchange happens. Scan speed, 8.4 s per frame. No postprocessing was done. (B) Four frames showing both direct exchange (frames 1 and 2) and SW transition (frames 2 to 4). Scale bars, 2 Å. Scan speed, 0.07 s per frame. A median filter



with a 2 pixel \times 2 pixel kernel was applied for clarity. The SW transition was captured during EELS acquisition in small subscan windows to enhance the signal-to-noise ratio of the spectra used to identify the dopants and to achieve faster scanning rate frames that can better capture atomic dynamics. (C) Neighboring C atom knocked out by the electron beam, turning a threefoldcoordinated P into fourfold-coordinated P. Scan speed, 8 s per frame. No postprocessing was done. (D) P dopant being replaced by a C atom. Scan speed, 4 s per frame. The different image color codings represent different categories: gray represents atom-conserving process, and magenta represents atomnonconserving process. Blue and red dashed circles in (A) and (B) represent the inequivalent lattice sites of graphene, and the green dashed circles in (C) and (D) indicate the location of the atom that has not been conserved. (E and F) Intentional control over the P direct exchange. The yellow crosses indicate the location where the electron beam was parked for 10 s to purposefully move the P atom by one lattice site. Green and blue dashed circles indicate the two nonequivalent lattices sites of graphene. Insets: The region of interest after applying a Gaussian filter. (G) A schematic plot of the control process, where the electron beam is represented by a green cone focused on the neighbor C atom. Credit: Science Advances (2019). advances.sciencemag.org/content/5/5/eaav2252

The ultimate degree of control for engineering would be the ability to create and manipulate materials at the most basic level, fabricating devices atom by atom with precise control.

Now, scientists at MIT, the University of Vienna, and several other institutions have taken a step in that direction, developing a method that can reposition atoms with a highly focused electron <u>beam</u> and control their exact location and bonding orientation. The finding could ultimately lead to new ways of making quantum computing devices or sensors, and usher in a new age of "atomic engineering," they say.

The advance is described today in the journal *Science Advances*, in a paper by MIT professor of nuclear science and engineering Ju Li,



graduate student Cong Su, Professor Toma Susi of the University of Vienna, and 13 others at MIT, the University of Vienna, Oak Ridge National Laboratory, and in China, Ecuador, and Denmark.

"We're using a lot of the tools of nanotechnology," explains Li, who holds a joint appointment in materials science and engineering. But in the new research, those tools are being used to control processes that are yet an order of magnitude smaller. "The goal is to control one to a few hundred atoms, to control their positions, control their charge state, and control their electronic and nuclear spin states," he says.

While others have previously manipulated the positions of individual atoms, even creating a neat circle of atoms on a surface, that process involved picking up <u>individual atoms</u> on the needle-like tip of a scanning tunneling microscope and then dropping them in position, a relatively slow mechanical process. The new process manipulates atoms using a relativistic electron beam in a scanning transmission electron microscope (STEM), so it can be fully electronically controlled by magnetic lenses and requires no mechanical moving parts. That makes the process potentially much faster, and thus could lead to practical applications.

Using electronic controls and artificial intelligence, "we think we can eventually manipulate atoms at microsecond timescales," Li says. "That's many orders of magnitude faster than we can manipulate them now with mechanical probes. Also, it should be possible to have many electron beams working simultaneously on the same piece of material."

"This is an exciting new paradigm for atom manipulation," Susi says.

Computer chips are typically made by "doping" a silicon crystal with other atoms needed to confer specific electrical properties, thus creating "defects' in the material—regions that do not preserve the perfectly orderly crystalline structure of the silicon. But that process is scattershot,



Li explains, so there's no way of controlling with atomic precision where those dopant atoms go. The new system allows for exact positioning, he says.

The same electron beam can be used for knocking an atom both out of one position and into another, and then "reading" the new position to verify that the atom ended up where it was meant to, Li says. While the positioning is essentially determined by probabilities and is not 100 percent accurate, the ability to determine the actual position makes it possible to select out only those that ended up in the right configuration.

Atomic soccer

The power of the very narrowly focused electron beam, about as wide as an atom, knocks an atom out of its position, and by selecting the exact angle of the beam, the researchers can determine where it is most likely to end up. "We want to use the beam to knock out atoms and essentially to play atomic soccer," dribbling the atoms across the graphene field to their intended "goal" position, he says.

"Like soccer, it's not deterministic, but you can control the probabilities," he says. "Like soccer, you're always trying to move toward the goal."

In the team's experiments, they primarily used <u>phosphorus atoms</u>, a commonly used dopant, in a sheet of graphene, a two-dimensional sheet of carbon atoms arranged in a honeycomb pattern. The phosphorus atoms end up substituting for carbon atoms in parts of that pattern, thus altering the material's electronic, optical, and other properties in ways that can be predicted if the positions of those atoms are known.

Ultimately, the goal is to move multiple atoms in complex ways. "We hope to use the <u>electron beam</u> to basically move these dopants, so we



could make a pyramid, or some defect complex, where we can state precisely where each atom sits," Li says.

This is the first time electronically distinct dopant atoms have been manipulated in graphene. "Although we've worked with silicon impurities before, phosphorus is both potentially more interesting for its electrical and magnetic properties, but as we've now discovered, also behaves in surprisingly different ways. Each element may hold new surprises and possibilities," Susi adds.

The system requires <u>precise control</u> of the beam angle and energy. "Sometimes we have unwanted outcomes if we're not careful," he says. For example, sometimes a carbon atom that was intended to stay in position "just leaves," and sometimes the phosphorus atom gets locked into position in the lattice, and "then no matter how we change the beam angle, we cannot affect its position. We have to find another ball."

Theoretical framework

In addition to detailed experimental testing and observation of the effects of different angles and positions of the beams and graphene, the team also devised a theoretical basis to predict the effects, called primary knock-on space formalism, that tracks the momentum of the "soccer ball." "We did these experiments and also gave a theoretical framework on how to control this process," Li says.

The cascade of effects that results from the initial beam takes place over multiple time scales, Li says, which made the observations and analysis tricky to carry out. The actual initial collision of the relativistic electron (moving at about 45 percent of the speed of light) with an atom takes place on a scale of zeptoseconds—trillionths of a billionth of a second—but the resulting movement and collisions of atoms in the lattice unfolds over time scales of picoseconds or longer—billions of



times longer.

Dopant atoms such as phosphorus have a nonzero nuclear spin, which is a key property needed for quantum-based devices because that spin state is easily affected by elements of its environment such as magnetic fields. So the ability to place these <u>atoms</u> precisely, in terms of both position and bonding, could be a key step toward developing quantum information processing or sensing devices, Li says.

More information: "Engineering single-atom dynamics with electron irradiation" *Science Advances* (2019). advances.sciencemag.org/content/5/5/eaav2252

This story is republished courtesy of MIT News (web.mit.edu/newsoffice/), a popular site that covers news about MIT research, innovation and teaching.

Provided by Massachusetts Institute of Technology

Citation: Manipulating atoms one at a time with an electron beam (2019, May 17) retrieved 2 May 2024 from <u>https://phys.org/news/2019-05-atoms-electron.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.