

Running may have made dinosaurs' wings flap before they evolved to fly

May 2 2019



Caudipteryx robot for testing passive flapping flight. Credit: Talori et al.

Before they evolved the ability to fly, two-legged dinosaurs may have



begun to flap their wings as a passive effect of running along the ground, according to new research by Jing-Shan Zhao of Tsinghua University, Beijing, and his colleagues.

The findings, published in *PLOS Computational Biology*, provide new insights into the origin of avian flight, which has been a point of debate since the 1861 discovery of *Archaeopteryx*. While a gliding type of flight appears to have matured earlier in <u>evolutionary history</u>, increasing evidence suggests that active flapping flight may have arisen without an intermediate gliding phase.

To examine this key point in evolutionary history, Zhao and his colleagues studied *Caudipteryx*, the most primitive, non-flying dinosaur known to have had feathered "proto-wings." This bipedal animal would have weighed around 5 kilograms and ran up to 8 meters per second.

First, the researchers used a mathematical approach called modal effective mass theory to analyze the mechanical effects of running on various parts of *Caudipteryx*'s body. These calculations revealed that running speeds between about 2.5 to 5.8 meters per second would have created forced vibrations that caused the dinosaur's wings to flap.

Real-world experiments provided additional support for these calculations. The scientists built a life-size robot of *Caudipteryx* that could run at different speeds, and confirmed that running caused a flapping motion of the wings. They also fitted a young ostrich with artificial wings and found that running indeed caused the wings to flap, with longer and larger wings providing a greater lift force.

"Our work shows that the motion of flapping feathered wings was developed passively and naturally as the dinosaur ran on the ground," Zhao says. "Although this flapping motion could not lift the dinosaur into the air at that time, the motion of flapping wings may have



developed earlier than gliding."

Zhao says that the next step for this research is to analyze the lift and thrust of *Caudipteryx*'s feathered wings during the passive flapping process.

More information: dentification of avian flapping motion from nonvolant winged dinosaurs based on modal effective mass analysis. *PLoS Comput Biol* 15(5): e1006846. <u>DOI: 10.1371/journal.pcbi.1006846</u>

Provided by Public Library of Science

Citation: Running may have made dinosaurs' wings flap before they evolved to fly (2019, May 2) retrieved 4 May 2024 from <u>https://phys.org/news/2019-05-dinosaurs-wings-evolved.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.