

The global invasion routes of the red swamp crayfish

May 16 2019



Red swamp crayfish specimens. Credit: Miguel Clavero


Spanish National Research Council (CSIC) researchers have

reconstructed the invasion routes followed by the red swamp crayfish during its human-driven expansion based on the analysis of a mitochondrial gene (COI), which was sequenced from 1,412 crayfishes from 122 populations across the Northern Hemisphere.

Invasion routes

The article describes how different invasion scenarios have produced different [genetic patterns](#) among invasive populations. "For example, in the US there are two main invasion routes: west- and east-wards from the native area. The invasive populations in the west are genetically more diverse, because they have received more introductions, which probably involved more specimens of crayfish, starting in the 1920s," explains Francisco J. Oficialdegui, CSIC researcher at the Doñana Biological Station.

The genetic results show that western US (California), itself an invaded area, was the source of the crayfish populations established in Hawaii and a probable source of the crayfish introduced to Japan, and from there to China, in the late 1920s. The low genetic diversity of all red [swamp](#) crayfish populations studied in Asia supports documentary evidence that a small group of some 20 individuals may have been the origin of the Japanese and Chinese red swamp crayfish populations which now number into the millions.



Invasion routes. Credit: Francisco J. Oficialdegui

Red swamp crayfish in Spain and Europe

The red swamp crayfish was introduced twice from Louisiana to southwestern Spain in 1973 (near the city of Badajoz) and 1974 (in the marshes of the Guadalquivir River). These introductions were promoted by the aristocrat Andrés Salvador de Habsburgo-Lorena. Until now, it has been assumed that these introductions were the sole origin of all red swamp crayfish populations established across Europe, but the new study finds evidence of a separate later introduction.

"The large number of individuals involved in the two introduction events (around 500 in Badajoz and 6,000 in the Guadalquivir marshes) has led to the high genetic diversity levels we observed in Iberian populations, although diversity values tend to be lower as populations are further away from the introduction foci. However, in our study we also

unexpectedly detected a genetic profile in central-western Europe that is not present in the Iberian Peninsula, a finding that suggests that additional unrecorded introductions of the red swamp [crayfish](#) into Europe may have occurred, either from the US or from other invaded territories," adds Oficialdegui.

More information: Francisco J. Oficialdegui et al, Unravelling the global invasion routes of a worldwide invader, the red swamp crayfish (*Procambarus clarkii*), *Freshwater Biology* (2019). [DOI: 10.1111/fwb.13312](#)

Provided by Spanish National Research Council (CSIC)

Citation: The global invasion routes of the red swamp crayfish (2019, May 16) retrieved 22 May 2024 from <https://phys.org/news/2019-05-global-invasion-routes-red-swamp.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--