

Turning off growth to make flowers grow

May 14 2019



Proper flower growth depends on the activation and suppression of floral stem cells (left). KNUCKLES and WUSCHEL expression are seen to colocalize in the nucleus to suppress floral stem cell activation (right). Credit: Toshiro Ito

The beautiful colors and smells of flowers serve a much greater purpose than just decoration. Flowers contain the plant's reproductive organs, and those same colors and smells that make a room beautiful also attract bees and other animals for pollination. Floral stem cells are crucial for the growth of the flower and its organs. That growth must eventually terminate for the flower to fully develop and set seeds. A new study led by scientists at the Nara Institute of Science and Technology (NAIST)



and seen in *The Plant Cell* shows that the transcription factor KNUCKLES is a key regulator of this stem cell arrest by initiating a serious of epigenetic events to repress the stem cell determinant WUSCHEL.

"Floral stem cell activity vanishes when WUSHCEL is suppressed and silenced through changes in its chromatin state. What we did not know was how this change begins and how it is sustained," explains NAIST Professor Toshiro Ito, who led the study.

Ito's team looked at the activation and suppression of floral stem cells from Arabidopsis. Stem cell activation was marked by a clear expression of WUSCHEL, but that changed when the <u>cells</u> also began to express KNUCKLES, which bound to the WUSCHEL locus and led to WUSCHEL's expression almost halving four hours later.

Then, at eight to 12 hours after the KNUCKLES expression, the group found that the WUSCHEL locus showed signs of H3K27me3 histone methylation, a marker of sustained gene suppression.

The question Ito wanted to answer was what were the events that took place from the KNUCKLES binding to the WUSCHEL locus to the H3K27me3 histone methylation that could terminate the stem cell activation.

"H3K27me3 is catalyzed by Polycomb Group complexes, but nothing is known about how the complexes are recruited to the WUSCHEL locus," says Ito.

The researchers discovered that KNUCKLES binding to WUSCHEL jettisoned SPLAYED, a chromatin remodeling protein that activates WUSCHEL. This effect leads to rapid transcriptional repression of WUSCHEL, followed by the recruitment of Polycomb Group complex



to WUSCHEL, where it formed H3K27me3 marks on the chromatin to suppress gene expression.

"KNUCKLES binding was essential for the rapid removal of active H3K4me3 marks and the following deposition of repressive H3K27me3 marks," explains Ito.

The recruitment was done by KNUCKLES interacting with a specific component of the Polycomb Group complex known as FERTILIZATION-INDEPENDENT ENDOSPERM.

"Our study reveals the temporal steps from KNUCKLES binding to H3K27me marks that silence the WUSCHEL chromatin. Understanding how stem cell activation is terminated will assist in new food technologies," says Ito.

More information: Bo Sun et al, Integration of transcriptional repression and Polycomb-mediated silencing of WUSCHEL in floral meristems, *The Plant Cell* (2019). DOI: 10.1105/tpc.18.00450

Provided by Nara Institute of Science and Technology

Citation: Turning off growth to make flowers grow (2019, May 14) retrieved 11 July 2024 from <u>https://phys.org/news/2019-05-growth.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.