

This hawk likes crab for dinner

May 2 2019



In 2016, authors observed a red-shouldered hawk attack and dismember another crustacean, the ghost crab (*Ocypode quadrata*), on a coastal beach in South Carolina, USA. Credit: Julie Larsen Maher/WCS

Red-shouldered hawks (*Buteo lineatus*) feed primarily on mammals, amphibians, reptiles, and invertebrates, the majority of which are insects and crustaceans, with the latter represented to date only by crayfish.

In 2016, authors observed a red-shouldered hawk attack and dismember another crustacean, the ghost crab (*Ocypode quadrata*), on a coastal beach in South Carolina, USA.

This is the first report of a red-shouldered [hawk](#) attacking and presumably consuming any species of crab and the first report of probable ghost crab predation by a raptor in North America.

More information: "'Attack and probable predation on ghost crab (*Ocypode quadrata*) by red-shouldered hawk (*Buteo lineatus*)," *Conservation Science and Practice* (2019). [DOI: 10.1676/18-151](https://doi.org/10.1676/18-151)

Provided by Wildlife Conservation Society

Citation: This hawk likes crab for dinner (2019, May 2) retrieved 2 May 2024 from <https://phys.org/news/2019-05-hawk-crab-dinner.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--