

How to prevent mosquitofish from spreading in water ecosystems

May 27 2019



Its high fecundity, sexual precocity and tolerance to environmental pollution are factors that make its monitoring more difficult. Credit: Oriol Cano (University of Barcelona-IRBio)

Preventing the introduction of the mosquitofish and removing its population are the most effective actions to control the dispersal of this exotic fish in ponds and lakes, according to a study published in the journal *Science of the Total Environment*. Neither the presence of predators nor the degradation of the quality of water and natural habitat are a threat to this invasive species -from the Atlantic coast in North

America- which competes against and moves local species away.

The new study, carried out in [water ecosystems](#) close to urban environments of the Barcelona provincial area, is signed by the experts Oriol Cano Rocabayera, Adolfo de Sostoa, Lluís Coll and Alberto Maceda, from the Faculty of Biology and the Biodiversity Research Institute of the University of Barcelona (IRBio).

An exotic species threatening local fish

The mosquitofish (*Gambusia holbrooki*) is an edacious exotic fish regarded as one of the most dangerous invasive [species](#) at a global scale, according to the International Union for the Conservation of Nature (UICN). It was brought to Spain in 1921 for the biological control of mosquitoes—carrying diseases such as malaria—and is now included in the catalogs of exotic invasive species of the Ministry of Agriculture, Fisheries and Food and the autonomous communities.

This species is especially present in the southern-western areas of the peninsula, the Mediterranean coast and the Ebro basin. In general, it is found in habitats with warm waters, with a low- or no-flow and [shallow waters](#)—in general, wetlands and the shallow waters in rivers—and areas with agricultural activity or peri-urban areas affected by pollution. In the peninsular area, it is a competitive predator that has forced [local species](#) to move -such as the Spanish toothcarp (*Aphanius iberus*), the Valencia toothcarp (*Valencia hispanica*), the Andalusian toothcarp (*Aphanius baeticus*) and the three-spined stickleback (*Gasterosteus aculeatus*), apart from other amphibians and invertebrates.

Exotic mosquitofish: strategies to survive under adverse conditions

Its high fecundity, sexual precocity, tolerance to environmental pollution and competitive superiority are "factors that make its monitoring more difficult if the only strategies to preserve biodiversity in the [water](#) ecosystem are based on the improvement of water or the [natural habitat](#)," notes researcher Oriol Cano Rocabayera, first author of the article and member of the Department of Evolutionary Biology, Ecology and Environmental Sciences of the UB and IRBio.

"If some mosquitofish enter a new habitat—or when there are a few of them after a control action to remove the population—this [exotic species](#) is able to increase its fecundity rate to balance the population imbalance. However, if the population of mosquitofish is stable and abundant, the fecundity lowers but new mosquitofish are bigger and have more chances of surviving," says Cano.

Changes in water regime-building of dams, canalizations, etc.—favor the presence of certain exotic species -like the mosquitofish—that prefer water with low flow. In general, the restoration of the hydrological regime has the best strategy to control invasive species.

"However, preventing the arrival of these and removing the populations of mosquitofish in the available ecosystems are the most effective actions for their control. Nevertheless, their [small size](#), the wide range of tolerance, their diet and connectivity of water ecosystems makes this control and their removal very expensive and difficult."

The new study reveals that habitats with abundant water plants—for example, naturalized dams—and well-preserved environments improve the physical condition of mosquitofish. Therefore, the complexity of the habitat is a factor that brings more preys and shelter against the attack of the predators. "The [mosquitofish](#)'s survival in high salinity waters is difficult, and these waters are now the natural habitat where the Spanish toothcarp is limited to," comments Cano Rocabayera.

The danger of releasing exotic animals in the environment

The global distribution of this [invasive species](#)—found in all continents except Antarctica—is related to the areas where it was introduced a century ago due the biological control of mosquitoes that carried malaria. Therefore, the most recent outbreaks of tropical diseases transmitted via mosquitoes—Zika, Chikunguya and Pappataci fever—make it essential to increase the surveillance on the these exotic fish in the local fauna, especially in tropical regions with a high biodiversity.

According to the authors of the study, "We need to apply more effective control measures on the small water masses, that is, in habitats where the complete removal of the population can be guaranteed. Also, we need to raise awareness among people on the danger of releasing exotic animals in the natural environment."

More information: Oriol Cano-Rocabayera et al, Managing small, highly prolific invasive aquatic species: Exploring an ecosystem approach for the eastern mosquitofish (*Gambusia holbrooki*), *Science of The Total Environment* (2019). [DOI: 10.1016/j.scitotenv.2019.02.460](https://doi.org/10.1016/j.scitotenv.2019.02.460)

Provided by University of Barcelona

Citation: How to prevent mosquitofish from spreading in water ecosystems (2019, May 27) retrieved 3 May 2024 from <https://phys.org/news/2019-05-mosquitofish-ecosystems.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.