

# From sharks in seagrass to manatees in mangroves, we've found large marine species in some surprising places

May 22 2019, by Michael Sievers, Rod Connolly And Tom Rayner

---



[https://images.theconversation.com/files/275559/original/file-20190521-23823-1ket9ng.jpg?ixlib=rb-1.1.0&q=45&auto=format&w=600&h=200&fit=crop&dp\\_r=1](https://images.theconversation.com/files/275559/original/file-20190521-23823-1ket9ng.jpg?ixlib=rb-1.1.0&q=45&auto=format&w=600&h=200&fit=crop&dp_r=1)

When we think of mangrove forests, seagrass meadows and saltmarshes, we don't immediately think of shark habitats. But the first [global review](#) of links between large marine animals (megafauna) and coastal wetlands is challenging this view—and how we might respond to the biodiversity crisis.

Mangrove forests, [seagrass meadows](#) and saltmarshes support rich [biodiversity](#), underpin the livelihoods of more than a billion people

worldwide, store carbon, and protect us from extreme weather events.

We know marine [megafauna](#) also use these habitats to live, feed and breed. Green turtles and manatees, for instance, are known to eat seagrass, and dolphins hunt in mangroves.

But new associations are also being discovered. The bonnethead shark—a close relative of hammerheads—was recently found to eat and digest seagrass.

The problem is that we're losing these important places. And until now, we've underestimated how important they are for large, charismatic and ecologically important marine animals.


## Counting wetland megafauna

Today our review of the connections between marine megafauna and vegetated coastal wetlands was [published in the journal \*Trends in Ecology and Evolution\*](#). As it turns out, far more megafauna [species](#) use coastal wetlands than we thought.

Before our review, the number of marine megafauna species known to use these habitats was 110, according to the [International Union for Conservation of Nature \(IUCN\) Red List](#), which assesses species' conservation status.

## Species that use vegetated coastal wetlands

Proportion of overall species group



Credit: The Conversation

We identified another 64 species from 340 published studies, bringing the total number to 174 species. This means 13% of all marine megafauna use vegetated coastal wetlands.

We predominantly documented these habitat associations by electronic tracking, direct observation or from analysing stomach contents or chemical tracers in animal tissues.

Less commonly, acoustic recordings and [animal-borne video](#) studies—strapping a camera on the back of turtle, for instance—were

used.

## Deepening our understanding of how species use their habitats

In recent weeks, the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES) released a damning [assessment](#) of humanity's stewardship of the natural world. Up to 1 million species were reported to be facing extinction within decades.

We need to dramatically change how we relate to and engage with species and their habitats, if we are to fix this problem.

But the question is, how can we make [global change](#) real, relevant and feasible at local and regional scales? And, as the international community rises to this challenge, what information is needed to support such efforts?

Our study suggests a critical first step to addressing the global [biodiversity crisis](#) is to deepen our understanding of links between species and their habitats. We also need to elevate how the evidence is used to both assess extinction risk and prioritise, plan and deliver conservation actions.

More than [half of all coastal wetlands have been lost globally](#) and the rest are at risk from a range of serious threats, including deforestation. There is an urgent need to limit and reverse the loss of [coastal wetlands](#) to stop biodiversity loss, protect communities and tackle climate change.

Targeting places where high rates of mangrove loss intersect with threatened megafauna could lead to more efficient and effective conservation outcomes. Southeast Asia, Mexico and northern Brazil are

[such places.](#)

In Southeast Asia, for example, the world's largest mangrove forest is losing trees at a rate far exceeding global averages, largely due to [aquaculture and agriculture](#). This is threatening the critically endangered green sawfish, which relies on these mangrove habitats.



Credit: Mile Ribeiro from Pexels

### **Habitats should always be considered in assessments**

The IUCN Red List assesses the extinction risk for almost 100,000 species. It provides comprehensive information on global conservation


statuses, combining information on population sizes, trends and threats.

The wealth of data collected during species' assessments, including habitat associations of threatened species, is one of the Red List's most valuable features.

But our study shows many known associations are yet to be included. And for more than half of the assessments for marine megafauna, habitat change is yet to be listed as a threat.

This is concerning because assessments that overlook habitat associations or lack sufficient detail, may not allow conservation resources be directed at the most effective recovery measures.

But it's also important to note habitat associations have varying strengths and degrees of supporting evidence. For example, a population of animals shown to consume substantial amounts of seagrass is clearly a stronger ecological link than an individual simply being observed above seagrass.



‘Proportion species’ refers to all species within key taxonomic groups that are associated with coastal wetlands. Author supplied

In our paper, we propose a simple framework to address these issues, by clarifying habitat associations in conservation assessments. Ideally, these assessments would include the following:

- list all habitat types the species is known to associate with
- indicate the type of association (occurrence, grazing, foraging or breeding)

- cite the source of supporting evidence
- provide an estimate of the level of habitat dependence.

## Data for decision making

Habitat loss is accelerating a global extinction crisis, but the importance of coastal habitats to marine megafauna has been significantly undervalued in assessments of extinction risk.

We need to strive to protect remaining coastal wetland habitats, not only for their ecological role, but also for their economic, social and cultural values to humans. We can do this by strengthening how we use existing scientific data on [habitat](#) associations in species assessments and conservation planning.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: From sharks in seagrass to manatees in mangroves, we've found large marine species in some surprising places (2019, May 22) retrieved 28 June 2024 from <https://phys.org/news/2019-05-sharks-seagrass-manatees-mangroves-weve.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.