


Astronomers see 'warm' glow of Uranus' rings

June 20 2019, by Robert Sanders



Composite image of Uranus's atmosphere and rings at radio wavelengths, taken with the ALMA array in December 2017. The image shows thermal emission, or heat, from the rings of Uranus for the first time, enabling scientists to determine their temperature: a frigid 77 Kelvin (-320 F). Dark bands in Uranus's atmosphere at these wavelengths show the presence of molecules that absorb radio waves, in particular hydrogen sulfide gas. Bright regions like the north polar spot (yellow spot at right, because Uranus is tipped on its side) contain very few of these molecules. Credit: UC Berkeley image by Edward Molter and Imke de Pater

The rings of Uranus are invisible to all but the largest telescopes—they weren't even discovered until 1977—but they're surprisingly bright in new heat images of the planet taken by two large telescopes in the high deserts of Chile.

The thermal glow gives astronomers another window onto the rings, which have been seen only because they reflect a little light in the visible, or optical, range and in the near-infrared. The new images taken by the Atacama Large Millimeter/submillimeter Array (ALMA) and the Very Large Telescope (VLT) allowed the team for the first time to measure the temperature of the rings: a cool 77 Kelvin, or 77 degrees above absolute zero—the boiling temperature of liquid nitrogen and equivalent to 320 degrees below zero Fahrenheit.

The observations also confirm that Uranus' brightest and densest ring, called the epsilon ring, differs from the other known ring systems within our solar system, in particular the spectacularly beautiful rings of Saturn.


"Saturn's mainly icy rings are broad, bright and have a range of particle sizes, from micron-sized dust in the innermost D ring, to tens of meters in size in the main rings," said Imke de Pater, a UC Berkeley professor

of astronomy. "The small end is missing in the main rings of Uranus; the brightest ring, epsilon, is composed of golf ball-sized and larger rocks."

By comparison, Jupiter's rings contain mostly small, micron-sized particles (a micron is a thousandth of a millimeter). Neptune's rings are also mostly dust, and even Uranus has broad sheets of dust between its narrow main rings.

"We already know that the epsilon ring is a bit weird, because we don't see the smaller stuff," said graduate student Edward Molter. "Something has been sweeping the smaller stuff out, or it's all glomming together. We just don't know. This is a step toward understanding their composition and whether all of the rings came from the same source material, or are different for each ring."

Rings could be former asteroids captured by the planet's gravity, remnants of moons that crashed into one another and shattered, the remains of moons torn apart when they got too close to Uranus, or debris remaining from the time of formation 4.5 billion years ago.



Near-infrared image of the Uranian ring system taken with the adaptive optics system on the 10-meter Keck telescope in Hawaii in July 2004. The image shows reflected sunlight. In between the main rings, which are composed of centimeter-sized or larger particles, sheets of dust can be seen. The epsilon ring seen in new thermal images is at the bottom. Credit: UC Berkeley image by Imke de Pater, Seran Gibbard and Heidi Hammel, 2006

The new data were published this week in *The Astronomical Journal*. De Pater and Molter led the ALMA observations, while Michael Roman and Leigh Fletcher from the University of Leicester in the United Kingdom led the VLT observations.

"The rings of Uranus are compositionally different from Saturn's main ring, in the sense that in optical and infrared, the albedo is much lower:

they are really dark, like charcoal," Molter said. "They are also extremely narrow compared to the rings of Saturn. The widest, the epsilon [ring](#), varies from 20 to 100 kilometers wide, whereas Saturn's are 100's or tens of thousands of kilometers wide."

The lack of dust-sized particles in Uranus' main rings was first noted when Voyager 2 flew by the planet in 1986 and photographed them. The spacecraft was unable to measure the temperature of the rings, however.

To date, astronomers have counted a total of 13 rings around the planet, with some bands of dust between the rings. The rings differ in other ways from those of Saturn.

"It's cool that we can even do this with the instruments we have," he said. "I was just trying to image the planet as best I could and I saw the rings. It was amazing."

Both the VLT and ALMA observations were designed to explore the temperature structure of Uranus' atmosphere, with VLT probing shorter wavelengths than ALMA.

"We were astonished to see the rings jump out clearly when we reduced the data for the first time," Fletcher said.

This presents an exciting opportunity for the upcoming James Webb Space Telescope, which will be able provide vastly improved spectroscopic constraints on the Uranian rings in the coming decade.

More information: Thermal Emission from the Uranian Ring System, arXiv:1905.12566 [astro-ph.EP] arxiv.org/abs/1905.12566

Provided by University of California - Berkeley

Citation: Astronomers see 'warm' glow of Uranus' rings (2019, June 20) retrieved 20 June 2024 from <https://phys.org/news/2019-06-astronomers-uranus.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.