


The atmosphere of a new ultra hot Jupiter is analyzed

June 11 2019



Credit: Instituto de Astrofísica de Canarias

The combination of observations made with the CARMENES spectrograph on the 3.5m telescope at Calar Alto Observatory (Almería), and the HARPS-N spectrograph on the National Galileo Telescope (TNG) at the Roque de los Muchachos Observatory (Garafía, La Palma) has enabled a team from the Instituto de Astrofísica de Canarias (IAC) to reveal new details about this extrasolar planet, which has a surface

temperature of around 2000 K.

MASCARA-2B/KELT-20b is an ultra hot Jupiter. It belongs to a new group of exoplanets, the hottest known until now, which can reach temperatures at the surface of over 2,000 K. The reason for its high temperature is the proximity of its orbit around its [host star](#), causing it to receive a large flux of radiation in the upper layers of its atmosphere.

The team, led by IAC researcher Núria Casasayas, which had already made initial measurements of the atmosphere during 2018, observed the planet during four transits. They used the HARP-N spectrograph on the National Galileo Telescope (TNG) at the Roque de los Muchachos Observatory (Garafía, La Palma) and the CARMENES spectrograph on the 3.5 m telescope of the Calar Alto Observatory (Almería).

"The two instruments sample slightly different wavelength regions, which allows us to sample a wider spectral range," explains Casasayas. She adds: "We have been able to detect hydrogen beta, singly ionized iron and magnesium with data from HARPS-N, while the presence of ionized calcium was detected only by using CARMENES. Neutral sodium and hydrogen alpha are detected with both instruments."

The study of exoplanetary atmospheres has become front line research in recent years. Instruments that perform high-resolution spectroscopy allow us not only to discover the atmospheric composition of planets outside the Solar System, but also to measure other important parameters such as the temperatures of the layers where their constituents are found, and other parameters of the dynamics of the atmosphere.

More information: N. Casasayas-Barris, et al. Atmospheric characterization of the ultra-hot Jupiter MASCARA-2b/KELT-20b. arXiv:1905.12491v1 [astro-ph.EP]. arxiv.org/abs/1905.12491

Provided by Instituto de Astrofísica de Canarias

Citation: The atmosphere of a new ultra hot Jupiter is analyzed (2019, June 11) retrieved 30 April 2024 from <https://phys.org/news/2019-06-atmosphere-ultra-hot-jupiter.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.