

Climate change may be putting beluga whales out of their depths

June 11 2019



Beluga whale. Credit: Wikipedia

An international team of researchers has found that the physical condition of beluga whales affects their capacity to store oxygen in their blood and muscle tissues, likely impacting their ability to dive. In a paper published in the *Journal of Experimental Biology*, concentrations of muscle myoglobin and blood hemoglobin, proteins responsible for the storage and delivery of oxygen, were found to be 12 and 27 percent higher, respectively, in beluga whales in peak physical condition compared to those of lower condition.

"We estimate that these differences equate to at least a three minute reduction in maximum dive time of the whales with the lowest [body condition](#)," says lead author Dr. Emily Choy (Ph.D./17), a University of Manitoba alumna, now a researcher at McGill University.

The study started in response to a 2014 report that found inferred growth rates of Beaufort Sea [beluga whales](#) had declined over the past 20 years, which were hypothesized to be the result of environmental change. As diving depth in toothed whales is related to body size, the team wondered if diving ability might be compromised by the smaller size of the whales.

However, the researchers found that body condition and not [body mass](#) was the most important factor predicting blood and muscle oxygen storage capacity. Previous work by Choy indicated that the northward migration of less energy dense prey species was driving the reductions in body condition.

"Considering that the peak biomass of Arctic cod, the beluga's main prey, is found between depths of 350 to 500 metres, and that the largest and most energy rich cod are found at the deepest depths, declines in body condition may affect the ability of the whales to reach their preferred prey," says Choy.



Dr. Emily Choy (PhD/17) says climate change is affecting whales' ability to dive for food. Credit: University of Manitoba

"The relationship between [physical condition](#) and oxygen storage capacity may be a vicious cycle in beluga whales, in which environmental changes resulting in decreased condition of the whales, affects their ability to forage, leading to further reductions in their condition through the diminished consumption of their prey," adds Dr. Michael Berenbrink, a study collaborator at the University of Liverpool.

University of Manitoba biologist Dr. Kevin Campbell notes: "Equally

concerning is that results from our study suggest environmental changes affecting the physical condition of beluga whales may also affect their ability to evade predators and sea ice entrapments, thus leading to increased mortality risks."

Working in collaboration with Inuit communities in the Inuvialuit Settlement Region of the Northwest Territories, the team was able to collect samples from 77 subsistence hunt beluga whales.

"We are very grateful for our [community partners](#), and would like to thank the hunters for sharing their [whales](#), without which this study would not have been possible," says Choy. "It is important that we continue to work together to monitor the health and condition of northern [beluga](#) populations."

More information: Emily S. Choy et al. Body condition impacts blood and muscle oxygen storage capacity of free-living beluga whales (*Delphinapterus leucas*), *The Journal of Experimental Biology* (2019). [DOI: 10.1242/jeb.191916](https://doi.org/10.1242/jeb.191916)

Provided by University of Manitoba

Citation: Climate change may be putting beluga whales out of their depths (2019, June 11) retrieved 16 June 2024 from <https://phys.org/news/2019-06-climate-beluga-whales-depths.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--