

Honey bee colonies down by 16%

June 5 2019



Credit: CC0 Public Domain

The number of honey bee colonies fell by 16 percent in the winter of 2017-18, according to an international study led by the University of Strathclyde.

The survey of 25,363 beekeepers in 36 countries found that, out of

544,879 colonies being managed at the start of winter, 89124 were lost, through a combination of circumstances including various effects of weather conditions, unsolvable problems with a colony's queen, and natural disaster.

Portugal, Northern Ireland, Italy and England experienced losses above 25 percent, while Belarus, Israel and Serbia were among those with loss rates below 10 percent. There were also significant regional variations within some countries, including Germany, Sweden and Greece.

The total loss rate was down from 20.9 percent in 2016-17 but was still higher than the 2015-16 figure of 12.0 percent. The total loss rate for Scotland increased over these three years, from 18.0 percent to 20.4 percent to 23.7 percent.

In a departure from previous findings, beekeepers who moved their colonies in the foraging season, to access other forage or for pollination, faced fewer losses than those who kept their bees in the same place. Smaller-scale beekeeping operations also had higher losses than larger ventures.

The study, based on voluntarily submitted information, covered 33 countries in Europe—including the four nations of the UK—along with Algeria, Israel and Mexico.

It has been published in the *Journal of Apicultural Research* and was carried out by researchers in the colony loss monitoring group of the international honey bee research association COLOSS, which is based in the Institute of Bee Health at the University of Bern.

Dr. Alison Gray, a Lecturer in Strathclyde's Department of Mathematics & Statistics, led the study. She said: "Loss of honey bee colonies is a highly complex issue. It tends to be influenced less by overall climate

than by specific weather patterns or a natural disaster affecting the colony. We observe colonies in winter but what happens to the bees then can be partly determined by the conditions of the previous summer.

"Many are also lost when there are problems with a colony's queen—for example, if she goes missing or is not laying the fertilised eggs which go on to become worker bees. Most colonies are also under attack from Varroa mites, a parasitic mite.

"The impact of beekeepers migrating their colonies would be expected to be partly dependent on the distance traveled and the reasons for migration; this would be worth further investigation."

The study had a focus on sources of forage, plants which bees visit to collect nectar and pollen, in six categories: orchards; oilseed rape; maize; sunflower; heather and autumn forage crops. These were potentially useful food sources for bees, which could help to build up a [colony](#), but, by extending the active and brood-rearing season of the bees, forage which was available late in the season could also extend the [reproductive cycle](#) for Varroa mites, weakening the bee colonies and making winter losses more likely.

More information: Alison Gray et al. Loss rates of honey bee colonies during winter 2017/18 in 36 countries participating in the COLOSS survey, including effects of forage sources, *Journal of Apicultural Research* (2019). [DOI: 10.1080/00218839.2019.1615661](https://doi.org/10.1080/00218839.2019.1615661)

Provided by University of Strathclyde, Glasgow

Citation: Honey bee colonies down by 16% (2019, June 5) retrieved 13 June 2024 from <https://phys.org/news/2019-06-honey-bee-colonies.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.