


Detection of powerful winds driven by a supermassive black hole from La Palma

June 7 2019



Credit: Instituto de Astrofísica de Canarias

The supermassive black holes in the centres of many galaxies seem to have a major influence on their evolution. This happens during a phase in which the black hole is consuming the material of the galaxy in which it resides at a very high rate, growing in mass as it does so. During this phase, the galaxy has an active galactic nucleus (AGN).

The effect that this activity has on the host galaxy is known as AGN feedback, and one of its properties is [galactic winds](#): This is gas from the centre of the galaxy being driven out by the energy released by the active nucleus. These winds can reach velocities of up to thousands of kilometres per second, and in the most energetic AGNs, for example, the quasars, which can clean out the centres of the galaxies impeding the formation of new stars. It has been shown that the evolution of the star formation over cosmological timescales cannot be explained without the existence of a regulating mechanism.

To study these winds in quasars, the EMIR infrared spectrograph on the Gran Telescopio Canarias (GTC) was used. EMIR was developed in the Instituto de Astrofísica de Canarias and is designed to study the coolest and most distant objects in the universe by analysing infrared light. Since June 2016 this has been installed at a focus of the GTC, after going through an exhaustive test phase in the workshops of the Instrument Division of the IAC headquarters in La Laguna.

The data obtained since then has been used to produce several [scientific articles](#) of which the latest is a study of the obscured quasar J1509+0434, published today in the journal *Monthly Notices of the Royal Astronomical Society Letters* and produced by an international team led by IAC researcher Cristina Ramos Almeida. This quasar is in the [local universe](#) and is an analogue of the more distant and far more numerous quasars in which AGN feedback must be affecting the formation of new stars in a major way.

"EMIR has allowed us to study the winds of ionized and [molecular gas](#) from this quasar by using the infrared range. This analysis is very important because they don't always show similar properties, which tells us a great deal about how these winds are produced and how they affect their host galaxies," explains Ramos Almeida. The study of this and other local quasars will allow us to understand what was happening in

galaxies when they were younger and when they were forming their structures which we see today.

Based on the new data obtained with EMIR, the team has discovered that the ionized [wind](#) is faster than the molecular wind, reaching velocities of up to 1,200 km/s. However, it would be the molecular wind which is emptying the gas reservoirs of the galaxy (up to 176 solar masses per year). "New observations with ALMA will let us confirm this estimate," explained José Acosta Pulido, a researcher at the IAC and co-author of this study.

The next step is to observe a complete sample of obscured nearby [quasars](#) with EMIR to study their ionized and molecular winds. We also want to investigate the stellar populations of their host galaxies. This will allow us to confirm the effect of AGN feedback on the evolution of the [galaxies](#) directly.

More information: C Ramos Almeida et al. A near-infrared study of the multiphase outflow in the type-2 quasar J1509+0434, *Monthly Notices of the Royal Astronomical Society: Letters* (2019). [DOI: 10.1093/mnrasl/slz072](#)

Provided by Instituto de Astrofísica de Canarias

Citation: Detection of powerful winds driven by a supermassive black hole from La Palma (2019, June 7) retrieved 11 May 2024 from <https://phys.org/news/2019-06-powerful-driven-supermassive-black-hole.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.