

Plant seeding and panspermia

June 27 2019, by Kevin Hattori



Professor Hagai Perets. Credit: American Technion Society

The first detection of an interstellar asteroid/comet-like object visiting the Solar system two years ago has sparked ideas about the possibility of interstellar travel. New research from the Technion–Israel Institute of Technology suggests that such objects also raise far reaching implications about the origins of planets across the galaxy, and possibly

even the initial formation of the Solar system itself.

The asteroid/comet-like object named 'Oumuamua confirmed decades-old scientific expectations that suggested that the interstellar medium is full of loose chunks of rock flying around. Such debris were thought to be ejected from planetary systems in the aftermath of planet formation, when large planets formed and kicked out some of the leftover minor-planets and planetesimals still lying around. From time to time, some fraction of these ejected rocks can still encounter foreign stars. In fortunate instances, this phenomenon can be observed as it swooshes through the Solar system.

The Technion researchers, Evgeni Grishin, Hagai Perets and Yael Avni wondered what would have happened if these 'Oumuamua-like interstellar rocks were flying around 4.5 billion years ago, when our star was young and wild, and a gaseous disk was present instead of our planetary system. Their findings could be critical to answering some of the biggest puzzles regarding planet formation and the origin of planets in the Solar system

Forming Planets with Foreign (Planetesimal) Currency

Planets are formed in protoplanetary disks, mostly made of gas and dust. The dust grains are thought to grow into pebbles, coagulate into bigger planetesimals, and finally, form planets. Once the objects reach km-size, they can survive and eventually coagulate and accrete smaller rocks/pebbles as to form planetary embryos and full-fledged planets. The main obstacle for such growth appears to occur before km-size objects form, in the stage when smaller rock and pebbles initially form. Indeed, several culprits conspire to destroy pebbles and meter-sized boulders before they can ever grow into larger planetesimals. Such pebbles and

rocks move through the gaseous disk in which they are initially embedded, and experience a headwind that slows them down.

The continuous push of the headwind might eventually lead them to quickly spiral inward into the Sun and be destroyed. In addition, collisions between small pebbles can lead to their fragmentation into smaller pieces halting their growth into larger planetesimals. In other words, pebbles and small rocks encounter a so-called "meter-size barrier" in their ability to grow into even larger planetesimals.

Several models were suggested as to overcome the meter-size barrier, but these typically require fine-tuned conditions that are unlikely to exist in most planetary systems; nevertheless, it is common knowledge that most if not all stars host planetary systems. The question is then how this came to be.

In their recently published paper in the *Monthly Notices of the Royal Astronomical Society*, Grishin and collaborators showed that interstellar objects are the key. They suggested that most systems do not need to go through the difficult stage of forming km-size planetesimals. Instead, most systems can capture interstellar km-size planetesimals that were originally ejected from other planetary systems. But how can an object moving at tens of km per second velocity through a Solar system be captured? It turns out the answer is simple—the same headwind that drives small rocks to inspiral into their sun can slow down bigger, km-size interstellar planetesimals and thereby capture them into a newly formed protoplanetary disk.

In this way, even a single planetary system can eject km-size planetesimals that then serve as seeds for the formation of many new planetary systems. As a result, even a very small number of planetary systems can seed the formation of many other systems—all it requires is just a few lucky rare cases to begin the process, and then these systems

can spawn planetesimal "seeds" across the galaxy, which in turn can be captured into a newly forming [protoplanetary disks](#) and provide them the basic km-size building blocks needed for planetary growth.

Planet formation no longer occurs in isolation; no planetary system is an island, but rather the reservoir of ejected rogue interstellar planetesimals serves to continuously initiate the birth of new planetary systems. In turn, any newly formed [planetary systems](#) ejects its own rogue planetesimals and help rebuild the reservoir of interstellar planetesimal seeds. The question then becomes: what are the odds of capturing these planetesimals, and how many successful formations are required to populate the entire birth cluster with planetesimals?

Nature Vs. Nurture: Where You Live Matters

To estimate the odds of planetesimal seeding and its implications for planet formation, the researchers developed a mathematical and numerical model for capture probability, depending on the properties of the interstellar planetesimal population and the disc. They found capturing small pebbles is extremely efficient, and that capturing larger bodies is more challenging, but still reasonable.

In the dense regions of stellar clusters where tens, hundreds, or even thousands of stars are born and live in a small regions (the "Manhattan" of star formation), around 10^6 'Oumuamuas are captured in the birth cluster, and the largest body captured can be as large as ~ 10 km.

In the galaxy's countryside, the galactic field environment, capture is more challenging, but still around $\sim 10^3$ 'Oumuamuas can be captured, and bodies up to ~ 1 km are captured per system—enough to serve as the seed for planet formation in each system.

One is Enough, Planetesimals Bring Joy and Life

The researchers summarize that only a small fraction of the stars in a cluster (less than 1 percent) are required to form the primordial planetesimals, which eventually seed the entire birth cluster of ~1000 stars. Roughly similar numbers are expected also for field environments. Both estimates are conservative. The interstellar reservoir therefore works in tandem with the main planet formation models, providing the initial seeds for many of the planetesimal formation models.

Another interesting side aspect is that biologically active material, in the form of bacteria, can survive the tough interstellar environment if the rock in which it is embedded is large enough (larger than a few cm scale). Although only a minute fraction of ejected rocks might harbor these hardcore bacteria, a large number of such potentially biologically active rocks can be captured. This gas-assisted capture is far more efficient mechanism for widespread panspermia, and most systems have probably gained their first life building blocks from somewhere else.

More information: Evgeni Grishin et al. Planet seeding through gas-assisted capture of interstellar objects, *Monthly Notices of the Royal Astronomical Society* (2019). [DOI: 10.1093/mnras/stz1505](https://doi.org/10.1093/mnras/stz1505)

Provided by American Technion Society

Citation: Planet seeding and panspermia (2019, June 27) retrieved 7 September 2024 from <https://phys.org/news/2019-06-seeding-panspermia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.