

Small steps, big leaps – how marram grass builds dunes

June 14 2019



Sand couch (left) uses dispersed growth patterns to build low, wide dunes. Marram grass (right) catches more sand immediately around it, resulting in a higher dune. Location: De Hors, Texel. Credit: Valérie Reijers

The size and shape of dunes varies greatly around the world: in Europe they're tall and narrow, while in the US they're low and wide. A new study has found that this is partly because dunes are constructed by plants with different 'movement strategies' determining the shape of the dune. The study was carried out by ecologists from Radboud University, the Royal Netherlands Institute for Sea Research (NIOZ) and the

University of Groningen. The findings, which will be published in *Nature Communications* on 14 June, could be useful for coastal landscape rehabilitation projects.

The coastline is a difficult place for plants to survive, thanks to all the salt and crashing waves. Marram grasses have a special survival mechanism to cope with this: they catch drifting sand and build a dune, which enables them to eventually escape the effects of the sea.

But why is it that our European marram [grass](#) constructs 15-metre dunes, while its American counterpart doesn't get higher than 6 metres? That was the question that researchers, including Valérie Reijers, a Ph.D. candidate at Radboud University, wanted to answer. They looked at how European and American marram grasses grow along the coast of Schiermonnikoog and in North Carolina.

'Moving' plants create the landscape

"Marram grasses create underground rootstocks which can develop new shoots. European marram grass appears to have a specific strategy in which short rootstocks create new shoots close to the older one, but occasionally there will be a long underground runner. This striking pattern of movement has previously been identified in other organisms, including mussels, albatrosses, sharks, bacteria and even humans," says Valérie Reijers, lead author of the article.

When searching for food, many animals move in a random pattern in which small movements alternate with a significant relocation, known as Lévy movements. "It's a sort of natural law that has emerged in different organisms," says Reijers. "It's striking that these 'movements' apparently also exist in plants and are so significant to the way in which they construct their [landscape](#)."

The use of Lévy movement enables European marram grass to capture a lot of drifting sand, in the most efficient way, with a minimal investment in its underground root network. American marram grass captures more sand by growing in a dispersed way, but this requires much more energy because the underground rootstocks have to travel a much greater distance. This results in a lower and wider [dune](#).

Self-regenerating dunes

Coastal landscapes are very important—they provide a habitat for many plant and animal species, capture CO₂ in the soil and, perhaps most importantly of all, they slow or break waves during storm surges.

"Natural coastal landscapes such as salt marshes and dunes are declining around the world," says Tjisse van der Heide, a researcher at NIOZ and professor of coastal ecology at the University of Groningen.

"Attempts to restore these landscapes are often unsuccessful. Plants are often uprooted by waves before they can establish themselves and capture any silt or sand. Now that we understand the spatial organisation of coastal plants, the next step is to apply these findings to regeneration projects so that we can help [plants](#) get established," says van der Heide.

Provided by Radboud University

Citation: Small steps, big leaps – how marram grass builds dunes (2019, June 14) retrieved 10 June 2023 from <https://phys.org/news/2019-06-small-big-marram-grass-dunes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.