

Development of 3-D particle model for single particles in battery electrodes

July 8 2019





Electrochemical phenomena inside a single particle predicted using the model developed. a) Charged state; b) Electrical potential; c) Lithium concentration; d) Voltage. Credit: DGIST



A model with a 3-D observation of micrometer-sized particles in a cell has been developed. Through the analysis and research of micrometersized particles in a cell, this model is expected to enhance energy efficiency of cells.

DGIST announced that Professor Yong Min Lee's team in the Department of Energy Science & Engineering developed a "micron single particle electrochemical model" that can estimate the electrochemical properties of a single particle of <u>electrode</u> active materials in 3-D. The 3-D observations of the single particles of electrode active materials, which are difficult to identify in an experiment, are expected to be applied to research on electrochemical phenomena and particle designs that enhance cell efficiency.

Although a secondary cell is commonly used as the power source of electric vehicles, it is still not as efficient an internal combustion engine. While efficiency can be improved by increasing the <u>energy</u> density of the <u>cells</u>, R&D has not been actively carried out due to the limitations in precise <u>analysis</u> technology.

Professor Lee's team posited that the energy density of a cell can be enhanced through the design optimization of electrode active materials in a cell. Then they sought a way to examine the micrometer-sized single particles of electrode active materials and developed electrochemical model that can conduct 3-D analysis on the single particles.





Experiment result that examined electrochemical phenomena by changing the design of single particle. Credit: DGIST

Unlike the existing model that focused on the cell electrode, the model developed by Professor Yong Min Lee's team focused on the single



particles of active materials that compose the electrode. By doing so, the team took another step closer to fundamentally increase cell efficiency through accurate analysis of the properties and characteristics of 3-D single particles in a model. Since it can have 3-D analysis of particles, the model is especially expected to be applied widely in research to design the single particles of electrode active <u>materials</u> in a cell.

Regarding this research, Professor Yong Min Lee in the Department of Energy Science and Engineering said "Comparing to previous works, our model can look into what happens within a single particle. As a result, it provides an innovative way in designing micrometer-sized <u>particles</u>. Our next goal is to apply this <u>electrochemical</u> model to improve the cell efficiency of electric vehicles."

More information: Jihun Song et al, 3D electrochemical model for a Single Secondary Particle and its application for operando analysis, *Nano Energy* (2019). DOI: 10.1016/j.nanoen.2019.05.087

Provided by DGIST (Daegu Gyeongbuk Institute of Science and Technology)

Citation: Development of 3-D particle model for single particles in battery electrodes (2019, July 8) retrieved 2 May 2024 from https://phys.org/news/2019-07-d-particle-particles-battery-electrodes.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.