

# Instability in Antarctic ice projected to make sea level rise rapidly

July 8 2019

---



Part of Thwaites Glacier crumbles into the ocean. It is part of the normal life of a glacier, but the rate of ice flow into the ocean of some Antarctic glaciers has markedly accelerated, raising concerns. Credit: NASA/OIB Jeremy Harbeck

Images of vanishing Arctic ice and mountain glaciers are jarring, but their potential contributions to sea level rise are no match for Antarctica's, even if receding southern ice is less eye-catching. Now, a study says that instability hidden within Antarctic ice is likely to

accelerate its flow into the ocean and push sea level up at a more rapid pace than previously expected.

In the last six years, five closely observed Antarctic glaciers have doubled their rate of ice loss, according to the National Science Foundation. At least one, Thwaites Glacier, modeled for the new study, may be in danger of succumbing to this instability, a volatile process that pushes ice into the ocean fast.

How much ice the glacier will shed in coming 50 to 800 years can't exactly be projected due to unpredictable fluctuations in climate and the need for more data. But researchers at the Georgia Institute of Technology, NASA Jet Propulsion Laboratory, and the University of Washington have factored the instability into 500 ice flow simulations for Thwaites with refined calculations.

The scenarios diverged strongly from each other but together pointed to the eventual triggering of the instability, which will be described in the question and answer section below. Even if [global warming](#) were to later stop, the instability would keep pushing ice out to sea at an enormously accelerated rate over the coming centuries.

And this is if [ice melt](#) due to warming oceans does not get worse than it is today. The study went with present-day ice melt rates because the researchers were interested in the instability factor in itself.

## **Glacier tipping point**

"If you trigger this instability, you don't need to continue to force the [ice sheet](#) by cranking up temperatures. It will keep going by itself, and that's the worry," said Alex Robel, who led the study and is an assistant professor in Georgia Tech's School of Earth and Atmospheric Sciences. "Climate variations will still be important after that tipping point because

they will determine how fast the ice will move."

"After reaching the tipping point, Thwaites Glacier could lose all of its ice in a period of 150 years. That would make for a [sea level rise](#) of about half a meter (1.64 feet)," said NASA JPL scientist Helene Seroussi, who collaborated on the study. For comparison, current sea level is 20 cm (nearly 8 inches) above pre-global warming levels and is blamed for increased coastal flooding.



Thwaites Glacier's outer edge. As the glacier flows into the ocean, it becomes sea ice and drives up sea level. Thwaites Glacier ice is flowing particularly fast, and some researchers believe it may have already tipped into instability or be near that point, though this has not yet been established. Credit: NASA/James Yungel

The researchers published their study in the journal the *Proceedings of*

*the National Academy of Sciences* on Monday, July 8, 2019. The research was funded by the National Science Foundation and NASA.

The study also showed that the instability makes forecasting more uncertain, leading to the broad spread of scenarios. This is particularly relevant to the challenge of engineering against flood dangers.

"You want to engineer critical infrastructure to be resistant against the upper bound of potential sea level scenarios a hundred years from now," Robel said. "It can mean building your [water treatment plants](#) and nuclear reactors for the absolute worst-case scenario, which could be two or three feet of sea level rise from Thwaites Glacier alone, so it's a huge difference."


## Q&A

### **Why is Antarctic ice the big driver of sea level rise?**

Arctic sea ice is already floating in water. Readers will likely remember that 90% of an iceberg's mass is underwater and that when its ice melts, the volume shrinks, resulting in no change in sea level.

But when ice masses long supported by land, like [mountain glaciers](#), melt, the water that ends up in the ocean adds to sea level. Antarctica holds the most land-supported ice, even if much of that land is seabed holding up just part of the ice's mass, while water holds up part of it. Also, Antarctica is an ice leviathan.

"There's almost eight times as much ice in the Antarctic ice sheet as there is in the Greenland ice sheet and 50 times as much as in all the mountain glaciers in the world," Robel said.



Ice melt at the grounding line contributes to seawater and thus sea levels, but the larger effect is to send more ice above it out into the water, where it also drives up sea level. When sea bottom behind the grounding line, under the ice, slopes downward going inland, it exacerbates the process, which can become unstable, perpetually pushing ice out to sea. Credit: antarcticglaciers.org, Creative Commons non-commercial license

### What is that 'instability' underneath the ice?

The line between where the ice sheet rests on the seafloor and where it extends over water is called the grounding line. In spots where the bedrock underneath the ice behind the grounding line slopes down, deepening as it moves inland, the instability can kick in.

On deeper beds, ice moves faster because water is giving it a little more lift. Also, warmer ocean water hollows out the bottom of the ice, adding a little more water to the ocean. More importantly, the ice above the hollow loses land contact and flows faster out to sea.

"Once ice is past the grounding line and just over water, it's contributing to sea level because buoyancy is holding it up more than it was," Robel said. "Ice flows out into the floating ice shelf and melts or breaks off as icebergs."

"The process becomes self-perpetuating," Seroussi said, describing why it is called "instability."

## **How did the researchers integrate instability into sea level forecasting?**

The researchers borrowed math from statistical physics that calculate what random variables do to predictability in a physical system, like ice flow, acted upon by outside forces, like temperature changes. They applied the math to simulations of possible future fates of marine [glaciers](#) like Thwaites Glacier.

They made an added surprising discovery. Normally, when climate conditions fluctuate strongly, Antarctic ice evens out the effects. Ice flow may increase but gradually, not wildly, but the instability produced the opposite effect in the simulations.

"The system didn't damp out the fluctuations, it actually amplified them. It increased the chances of rapid ice loss," Robel said.

## **How rapid is 'rapid' sea level rise and when will we feel it?**

The study's time scale was centuries, as is common for sea level studies. In the simulations, Thwaites Glacier colossal ice loss kicked in after 600 years, but it could come sooner.

"It could happen in the next 200 to 600 years. It depends on the bedrock topography under the ice, and we don't know it in great detail yet," Seroussi said.

So far, Antarctica and Greenland have lost a small fraction of their ice, but already, shoreline infrastructures face challenges from increased tidal flooding and storm surges. Sea level is expected to rise by up to two feet by the end of this century.

For about 2,000 years until the late 1800s, sea level held steady, then it began climbing, according to the Smithsonian Institution. The annual rate of sea level rise has roughly doubled since 1990.

**More information:** Alexander A. Robel et al., "Marine ice sheet instability amplifies and skews uncertainty in projections of future sea-level rise," *PNAS* (2019).

[www.pnas.org/cgi/doi/10.1073/pnas.1904822116](http://www.pnas.org/cgi/doi/10.1073/pnas.1904822116)

Provided by Georgia Institute of Technology

Citation: Instability in Antarctic ice projected to make sea level rise rapidly (2019, July 8) retrieved 19 April 2024 from

<https://phys.org/news/2019-07-instability-antarctic-ice-sea-rapidly.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.