


# On the way to nanotheranostics

July 8 2019



Near-infrared excitation of functionalized bismuth ferrite nanoparticles results in harmonic emission which triggers the photorelease of caged molecular cargos, allowing decoupled imaging and controlled delivery applications. Credit: S. Gerber/EPFL

Theranostics is an emerging field of medicine whose name is a combination of "therapeutics" and "diagnostics." The idea behind theranostics is to combine drugs and/or techniques to simultaneously—or sequentially—diagnose and treat medical conditions, and also monitor

the response of the patient. This saves time and money, but can also bypass some of the undesirable biological effects that may arise when these strategies are employed separately.

Today, theranostics applications increasingly use nanoparticles that unite diagnostic molecules and drugs into a single agent. The nanoparticles act as carriers for molecular "cargo," e.g. a drug or a radioisotope to [cancer patients](#) undergoing radiotherapy, targeting specific biological pathways in the patient's body, while avoiding damage to healthy tissues.

Once at their target tissue, the nanoparticles produce diagnostic images and/or deliver their cargo. This is the cutting-edge technology of "nanotheranostics," which has become a major focus of research—albeit with many limitations to overcome.

Now, the lab of Sandrine Gerber at EPFL, working with physicists at the University of Geneva, have developed a new nanotheranostic system that overcomes several problems with previous approaches. The system uses "harmonic nanoparticles" (HNPs), a family of metal-oxide nanocrystals with exceptional optical properties, in particular their emission in response to excitation from ultraviolet to [infrared light](#), and their high photostability. It was this feature that brought HNPs into nanotheranostics, when scientists were trying to solve some problems with fluorescent probes.

"Most light-activated nanotheranostic systems need high-energy UV light to excite their photoresponsive scaffolds," says Gerber. "The problem is that this results in poor penetration depth and can damage living cells and tissues, which limits biomedical applications."

The new system that Gerber's group developed avoids these problems by using silica-coated bismuth-ferrite HNPs functionalized with light-responsive caged molecular cargos. These systems can be easily

activated with near-infrared light (wavelength 790 nanometers) and imaged at longer wavelength for both detection and drug release processes. Both these features render the system medically safe for patients.

Once light-triggered, the HNPs release their cargo—in this case, L-tryptophan, used as a model. The scientists monitored and quantified the release with a technique that combines [liquid chromatography](#) and [mass spectrometry](#), covering the imaging-diagnostic part of the nanotheranostic system.

The authors state that "this work is an important step in the development of nanocarrier platforms allowing decoupled imaging in tissue depth and on-demand release of therapeutics."

**More information:** Jérémy Vuilleumier et al. Two-photon triggered photorelease of caged compounds from multifunctional harmonic nanoparticles, *ACS Applied Materials & Interfaces* (2019). [DOI: 10.1021/acsami.9b07954](#)

Provided by Ecole Polytechnique Federale de Lausanne

Citation: On the way to nanotheranostics (2019, July 8) retrieved 25 April 2024 from <https://phys.org/news/2019-07-nanotheranostics.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.