


How nature builds hydrogen-producing enzymes

July 24 2019



Thomas Happe (left) and Oliver Lampret aim at understanding the workings of hydrogen producing enzymes in detail. Credit: RUB, Kramer

A team from Ruhr-Universität Bochum and the University of Oxford has discovered how hydrogen-producing enzymes, called hydrogenases,

are activated during their biosynthesis. They showed how the cofactor—part of the active centre and also the heart of the enzyme—is introduced inside.

Hydrogenases are of biotechnological interest as they are able to efficiently produce hydrogen. "In order to optimise them for an industrial application, we first need to understand the process of how the [protein](#) shell takes up and activates the chemical cofactor," says Professor Thomas Happe. A team led by Oliver Lampret and Thomas Happe from the Bochum-based Photobiotechnology Research Group published the results in the journal *Proceedings of the National Academy of Sciences*.

The researchers investigated the subgroup of [FeFe]-hydrogenases, which are the most efficient hydrogen producers. In nature, they can be found in green algae. Within their [protein scaffold](#), the enzymes have an active centre, the so-called H-cluster, where the hydrogen is produced. It is comprised of two structural elements: a cluster containing four iron and four sulphur atoms, and the catalytic cofactor, which consists of two iron and two sulphur atoms. "This cofactor is the linchpin of the [enzyme](#)," explains Oliver Lampret.

Final step of biosynthesis

In nature, the cofactor is subsequently incorporated into the enzyme following the biosynthesis of the protein scaffold—a highly complex process. Only then is the hydrogenase catalytically active. The researchers clarified the precise sequence of the process using protein engineering, protein film electrochemistry and infrared spectroscopy.

The team showed that the negatively charged cofactor is specifically transported through a positively charged maturation channel into the interior of the enzyme before it is firmly anchored into the protein shell.

Particularly flexible [structural elements](#) act as hinges here, ensuring that the protein folds differently and firmly envelopes and protects the integrated cofactor. The interaction of protein environment and cofactor is essential in order to stabilise the cofactor in its catalytic form.

"We assume that not only do [FeFe]-hydrogenases obtain their cofactor in this way but that the mechanism also occurs in other metalliferous enzymes," says Happe.

More information: Oliver Lampret et al. The final steps of [FeFe]-hydrogenase maturation, *Proceedings of the National Academy of Sciences* (2019). [DOI: 10.1073/pnas.1908121116](https://doi.org/10.1073/pnas.1908121116)

Provided by Ruhr-Universitaet-Bochum

Citation: How nature builds hydrogen-producing enzymes (2019, July 24) retrieved 2 May 2024 from <https://phys.org/news/2019-07-nature-hydrogen-producing-enzymes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.