


Physicists have let light through the plane of the world's thinnest semiconductor crystal

July 23 2019



As the scientists noted, the graph that shows the spatial distribution of the polarization of light turned out to be rather unusual -- it resembles a multi-colored marine rapan. Credit: *Nature Nanotechnology*

In every modern microcircuit hidden inside a laptop or smartphone, you can see transistors—small semiconductor devices that control the flow of electric current, i.e. the flow of electrons. If we replace electrons with photons (elementary particles of light), then scientists will have the prospect of creating new computing systems that can process massive information flows at a speed close to the speed of light. At present, it is photons that are considered the best for transmitting information in quantum computers. These are still hypothetical computers that live according to the laws of the quantum world and are able to solve some problems more efficiently than the most powerful supercomputers.

Although there are no fundamental limits for creating [quantum](#) computers, scientists still have not chosen what material platform will be the most convenient and effective for implementing the idea of a quantum computer. Superconducting circuits, cold atoms, ions, defects in diamond and other systems now compete for being one chosen for the future quantum computer. It has become possible to put forward the semiconductor platform and [two-dimensional crystals](#), specifically, thanks to scientists from: the University of Würzburg (Germany); the University of Southampton (United Kingdom); the University of Grenoble Alpes (France); the University of Arizona (USA); the Westlake university (China), the Ioffe Physical Technical Institute of the Russian Academy of Sciences; and St Petersburg University.

The physicists studied the propagation of light in a two-dimensional crystal layer of molybdenum diselenide (MoSe_2) which is only one atom thick—this is the thinnest semiconductor crystal in the world. The researchers found that the polarisation of light propagating in a superfine crystalline layer depends on the direction of light propagation. This phenomenon is due to the effects of spin-orbit interaction in the crystal. Interestingly, as the scientists noted, the graph that shows the spatial distribution of the polarisation of light turned out to be rather unusual—it resembles a multi-coloured marine rapana.

Ultrafine molybdenum diselenide crystals for experiments were synthesised in the laboratory of Professor Sven Höfling at the University of Würzburg. It is one of the best crystal growth laboratories in Europe. Measurements were carried out both in Würzburg and in St Petersburg under the supervision of Alexey Kavokin, professor at St Petersburg University. An important role in the development of the theoretical base was made by Mikhail Glazov. He is a corresponding member of the Russian Academy of Sciences, an employee of the Spin Optics Laboratory at St Petersburg University, and a leading research associate at the Ioffe Physical Technical Institute.

"I foresee that in the near future, two-dimensional monoatomic crystals will be used to transfer information in quantum devices," said Professor Alexey Kavokin, head of the Spin Optics Laboratory at St Petersburg University. "What classic computers and supercomputers take a very long time to do, a [quantum computing](#) device will do very quickly. Therein lies the great danger of quantum technologies—comparable to the danger of an atomic bomb. With their help it will be possible, for example, to hack banking protection systems very quickly. That is why today intensive work is under way, including the creation of means of protecting quantum devices: quantum cryptography. And our work contributes to semiconductor quantum technologies."

Additionally, as the scientist noted, the research was a major step forward in the study of light-induced (i.e. appearing in the presence of light) superconductivity. It is the phenomenon when the materials that allow electric current to pass through have zero resistance. At present, this state cannot be achieved at temperatures above minus 70 C. However, if the proper material is found, this discovery will make it possible to transfer electricity to any point on Earth without any loss, and to create a new generation of electric motors. It should be recalled that in March 2018, the research team of Alexey Kavokin predicted that structures containing superconducting metals, such as aluminium, can

help solve the problem. Nowadays, scientists at St Petersburg University are looking for a way to obtain experimental evidence of their theory.

More information: Nils Lundt et al, Optical valley Hall effect for highly valley-coherent exciton-polaritons in an atomically thin semiconductor, *Nature Nanotechnology* (2019). [DOI: 10.1038/s41565-019-0492-0](https://doi.org/10.1038/s41565-019-0492-0)

Provided by St. Petersburg State University

Citation: Physicists have let light through the plane of the world's thinnest semiconductor crystal (2019, July 23) retrieved 30 April 2024 from <https://phys.org/news/2019-07-physicists-plane-world-thinnest-semiconductor.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--