

# **Super salty, subzero Arctic water provides peek at possible life on other planets**

July 12 2019, by Hannah Hickey

---



Zac Cooper and Shelly Carpenter begin to drill below the Alaskan ice tunnel toward the cryopeg and its salty subzero water. Researchers are careful to sterilize their equipment to avoid introducing contamination from above ground. The most stringent of such techniques will be needed to sample for life on other planets. Credit: Go Iwahana/University of Alaska, Fairbanks


In recent years, the idea of life on other planets has become less far-fetched. NASA announced June 27 that it will send a vehicle to Saturn's icy moon Titan, a celestial body known to harbor surface lakes of methane and an ice-covered ocean of water, boosting its chance for supporting life.

On Earth, scientists are studying the most extreme environments to learn how life might exist under completely different settings, like on other planets. A University of Washington team has been studying the microbes found in "cryopegs," trapped layers of sediment with water so salty that it remains liquid at below-freezing temperatures, which may be similar to environments on Mars or other planetary bodies farther from the sun.

At the recent AbSciCon meeting in Bellevue, Washington, researchers presented DNA sequencing and related results to show that brine samples from an Alaskan cryopeg isolated for tens of thousands of years contain thriving bacterial communities. The lifeforms are similar to those found in floating sea ice and in saltwater that flows from glaciers, but display some unique patterns.

"We study really old seawater trapped inside of permafrost for up to 50,000 years, to see how those bacterial communities have evolved over time," said lead author Zachary Cooper, a UW doctoral student in oceanography.





A schematic of the study site, which consists of a tunnel, excavated from a massive ice formation in the permafrost, and accessed through a narrow vertical opening. Researchers then drill below the tunnel floor to reach the cryopeg layer with its saline liquid (bottom hatched area). Credit: Shelly Carpenter/University of Washington

Cryopegs were first discovered by geologists in Northern Alaska decades ago. This field site in Utqiagvik, formerly known as Barrow, was

excavated in the 1960s by the U.S. Army's Cold Regions Research and Engineering Laboratory to explore large wedges of freshwater ice that occur in the permafrost there. Subsurface brine was eventually collected from the site in the 2000s.

"The [extreme conditions](#) here are not just the below-zero temperatures, but also the very high salt concentrations," said Jody Deming, a UW professor of oceanography who studies microbial life in the Arctic Ocean. "One hundred and forty parts per thousand—14%—is a lot of salt. In canned goods that would stop microbes from doing anything. So there can be a preconceived notion that very high salt should not enable active life."

It's not fully known how cryopegs form. Scientists believe the layers might be former coastal lagoons stranded during the last ice age, when rain turned to snow and the [ocean](#) receded. Moisture evaporated from the abandoned seabed was then covered by permafrost, so the remaining briny water became trapped below a layer of frozen soil.



The research site about 1 mile outside of Utqiagvik, Alaska, appears at the surface as a box sitting on an expanse of white tundra. This is one of two cryopeg locations under study worldwide. It's not known how many of these features exist, but evidence suggests they're widespread in flat Arctic coastal regions. Credit: Zac Cooper/University of Washington







Zac Cooper taking notes inside the ice tunnel, with light from his headlamp. The team spends four- to eight-hour shifts inside the tunnel. One person gets the luxury of sitting on a bucket. Credit: Shelly Carpenter/University of Washington



The roof of the tunnel is covered in hoar frost, spiky ice crystals that form as moisture in the air solidifies in the minus 6 degrees C environment of the tunnel. The layers below are colder. Researchers leave presterilized pipes inserted in the floor for future access to the liquid layer below. Credit: Zac Cooper/University of Washington



To access the subsurface liquids, researchers climb about 12 feet down a ladder and then move carefully along a tunnel within the ice. The opening is just a single person wide and is not high enough to stand in, so researchers must crouch and work together to drill during the 4- to 8-hour shifts.

Deming describes it as "exhilarating" because of the possibility for discovery.

Samples collected in the spring of 2017 and 2018, geologically isolated for what researchers believe to be roughly 50,000 years, contain genes from healthy communities of bacteria along with their viruses.



Oceanography graduate student Zac Cooper climbs down an icy ladder into the tunnel in May 2018. Researchers are harnessed to a rope for safety. Credit: Shelly Carpenter/University of Washington

"We're just discovering that there's a very robust microbial community, coevolving with viruses, in these ancient buried brines," Cooper said. "We were quite startled at how dense the bacterial communities are."

The [extreme environments](#) on Earth may be similar to the oceans and ice of other planets, scientist believe.

"The dominant bacterium is Marinobacter," Deming said. "The name alone tells us that it came from the ocean—even though it has been in the dark, buried in frozen permafrost for a very long time, it originally came from the marine environment."

Mars harbored an ocean of water in the past, and our solar system contains at least a half-dozen oceans on other planets and [icy moons](#). Titan, the moon of Saturn that NASA will explore, is rich in various forms of ice. Studying life on Earth in frozen settings that may have similarities can prepare explorers for what kind of [life](#) to expect, and how to detect it.

Provided by University of Washington

Citation: Super salty, subzero Arctic water provides peek at possible life on other planets (2019, July 12) retrieved 3 May 2024 from <https://phys.org/news/2019-07-super-salty-subzero-arctic-peek.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is
---



provided for information purposes only.