

Rising tundra temperatures create worrying changes in microbial communities

July 8 2019



Researchers studied the impact of warming on microbial communities in a tundra area near Denali National Park in Alaska. Credit: Professor Ted Schuur, Northern Arizona University

Rising temperatures in the tundra of the Earth's northern latitudes could affect microbial communities in ways likely to increase their production of greenhouse gases methane and carbon dioxide, a new study of experimentally warmed Alaskan soil suggests.

About half of the world's total underground carbon is stored in the soils of these frigid, northern latitudes. That is more than twice the amount of carbon currently found in the atmosphere as [carbon dioxide](#), but until now most of it has been locked up in the very cold soil. The new study, which relied on metagenomics to analyze changes in the [microbial communities](#) being experimentally warmed, could heighten concerns about how the release of this carbon may exacerbate [climate change](#).

"We saw that microbial communities respond quite rapidly—within four or five years—to even modest levels of [warming](#)," said Kostas T. Konstantinidis, the paper's corresponding author and a professor in the School of Civil and Environmental Engineering and the School of Biological Sciences at the Georgia Institute of Technology.

"Microbial species and their genes involved in carbon dioxide and methane release increased their abundance in response to the warming treatment. We were surprised to see such a response to even mild warming."

The new study was supported by the U.S. Department of Energy and the National Science Foundation, and reported July 8 in the early edition of the journal *Proceedings of the National Academy of Sciences*. Researchers from the University of Oklahoma, Michigan State University and Northern Arizona University collaborated with Georgia Tech on the study.

The study provides quantitative information about how rapidly microbial communities responded to the warming at critical depths, and highlights

the dominant microbial metabolisms and groups of organisms that are responding to warming in the tundra. The work underscores the importance of accurately representing the role of soil microbes in climate models.



Test plots were used by researchers to study the effects of warming on microbial communities in the interior Alaskan landscape. Credit: Professor Ted Schuur, Northern Arizona University

The research began in September 2008 at a moist, acidic tundra area in the interior of Alaska near Denali National Park. Six experimental

blocks were created, and in each block, two snow fences were constructed about five meters apart in the winter to control snow cover. Thicker snow cover in the winter served as an insulator, creating slightly elevated temperatures—about 1.1 degrees Celsius (2 degrees Fahrenheit) in the experimental plots.

Other than the temperature difference, the soil conditions were similar in the experimental and control plots. Soil cores were taken from the experimental and control plots at two different depths at two different times: 1.5 years after the experiment began, and 4.5 years after the start.

Microbial DNA was extracted from the cores and sequenced using the Genomics Core at Georgia Tech.

"Our analysis of the resulting data showed which species were there, in what abundances, which species responded to warming and by how much—and what functions they possessed related to carbon use and release," said Eric R. Johnston, now a postdoctoral researcher at Oak Ridge National Laboratory, who conducted the study's analysis as a Georgia Tech Ph.D. student.

Cores from the experimental and control plots were compared to assess the effects of the warming. Cumulative ecosystem respiration was also sampled during the month following removal of the cores.

"The response we observed differed markedly between the two soil depths (15 to 25 centimeters and 45 to 55 centimeters) that were sampled for this study," said Johnston. "Specifically, at the upper boundary of the initial permafrost boundary layer—45 to 55 centimeters below the surface—the relative abundance of genes involved in methane production (methanogenesis) increased with warming, while genes involved in organic carbon respiration—the release of carbon dioxide—became more abundant at shallower depths."



Cumulative respiration from the tundra microbial communities was sampled during the month following removal of the soil cores. Credit: Professor Ted Schuur, Northern Arizona University

Measurement of the community respiration showed increases in the rate of carbon dioxide and methane release in the plots that were warmed. "Similar measurements have also shown that these gases are being released at a greater rate throughout the entire region in recent years as a result of climate warming," Johnston added.

The two soil depths correspond to an active layer near the surface that

freezes during the winter but thaws during warmer months, exposing the carbon. The deeper measurements examined soil just above the permafrost that thaws for only a brief time each year. These variations create fundamental differences in the biology and chemistry at the two depths.

"We expected to observe warming responses that differed between the two sampling depths," Johnston said. "Ongoing thaw of permafrost soil is being observed on the global scale, so we were particularly interested in evaluating microbiological responses to thawing permafrost."

The research highlights the importance of microbial communities in contributing atmospheric methane and carbon dioxide to climate change, Konstantinidis said.

"Because of the very large amount of [carbon](#) in these systems, as well as the rapid and clear response to warming found in this experiment and other studies, it is becoming increasingly clear that [soil](#) microbes—particularly those in the northern latitudes—and their activities need to be represented in climate models," he said. "Our work provides markers—species and genes—that can be used in this direction."

More information: Eric R. Johnston et al., "Responses of tundra soil microbial communities to half a decade of experimental warming at two critical depths," *PNAS* (2019).

www.pnas.org/cgi/doi/10.1073/pnas.1901307116

Provided by Georgia Institute of Technology

Citation: Rising tundra temperatures create worrying changes in microbial communities (2019,

July 8) retrieved 23 April 2024 from <https://phys.org/news/2019-07-tundra-temperatures-microbial.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.