

## Two advances in understanding the role of 'charge stripes' in superconducting mate

August 19 2019, by Ali Sundermier And Glennda Chu



This cutaway view shows stripes of higher and lower electron density -- 'charge stripes' -- within a copper-based superconducting material. Experiments with SLAC's X-ray laser directly observed how those stripes fluctuate when hit with a pulse of light, a step toward understanding how they interact with high-temperature superconductivity. Credit: Greg Stewart/SLAC National Accelerator Laboratory

High-temperature superconductors, which carry electricity with zero



resistance at much higher temperatures than conventional superconducting materials, have generated a lot of excitement since their discovery more than 30 years ago because of their potential for revolutionizing technologies such as maglev trains and long-distance power lines. But scientists still don't understand how they work.

One piece of the puzzle is the fact that <u>charge density waves</u>—static stripes of higher and lower electron density running through a material—have been found in one of the major families of <u>high-</u> <u>temperature superconductors</u>, the copper-based cuprates. But do these charge stripes enhance superconductivity, suppress it or play some other role?

In independent studies, two research teams report important advances in understanding how charge stripes might interact with superconductivity. Both studies were carried out with X-rays at the Department of Energy's SLAC National Accelerator Laboratory.

## **Exquisite detail**

In a paper published today in *Science Advances*, researchers from the University of Illinois at Urbana-Champaign (UIUC) used SLAC's Linac Coherent Light Source (LCLS) X-ray free-electron laser to observe fluctuations in charge density waves in a cuprate superconductor.

They disturbed the charge density waves with pulses from a conventional laser and then used RIXS, or resonant inelastic X-ray scattering, to watch the waves recover over a period of a few trillionths of a second. This recovery process behaved according to a universal dynamical scaling law: It was the same at all scales, much as a fractal pattern looks the same whether you zoom in or zoom out.

With LCLS, the scientists were able to measure, for the first time and in



exquisite detail, exactly how far and how fast the charge density waves fluctuated. To their surprise, the team discovered that the fluctuations were not like the ringing of a bell or the bouncing of a trampoline; instead, they were more like the slow diffusion of a syrup—a quantum analog of liquid crystal behavior, which had never been seen before in a solid.

"Our experiments at LCLS establish a new way to study fluctuations in charge density waves, which could lead to a new understanding of how high-temperature superconductors operate," says Matteo Mitrano, a postdoctoral researcher in professor Peter Abbamonte's group at UIUC.

This team also included researchers from Stanford University, the National Institute of Standards and Technology and Brookhaven National Laboratory.

## **Hidden arrangements**

Another study, reported last month in *Nature Communications*, used X-rays from SLAC'S Stanford Synchrotron Radiation Lightsource (SSRL) to discover two types of charge density wave arrangements, making a new link between these waves and high-temperature superconductivity.

Led by SLAC scientist Jun-Sik Lee, the research team used RSXS, or resonant soft X-ray scattering, to watch how temperature affected the charge <u>density</u> waves in a cuprate superconductor.

"This resolves a mismatch in data from previous experiments and charts a new course for fully mapping the behaviors of electrons in these exotic superconducting materials," Lee says.

"I believe that exploring new or hidden arrangements, as well as their intertwining phenomena, will contribute to our understanding of high-



temperature superconductivity in cuprates, which will inform researchers in their quest to design and develop new <u>superconductors</u> that work at warmer temperatures."

**More information:** J.-J. Wen et al. Observation of two types of chargedensity-wave orders in superconducting  $La_2-xSr_xCuO_4$ , *Nature Communications* (2019). DOI: 10.1038/s41467-019-11167-z

Matteo Mitrano et al. Ultrafast time-resolved x-ray scattering reveals diffusive charge order dynamics in La2–xBaxCuO4, *Science Advances* (2019). DOI: 10.1126/sciadv.aax3346

Provided by SLAC National Accelerator Laboratory

Citation: Two advances in understanding the role of 'charge stripes' in superconducting mate (2019, August 19) retrieved 4 May 2024 from <u>https://phys.org/news/2019-08-advances-role-stripes-superconducting.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.