

Research sheds new light on Antarctic control of global climate

August 28 2019



Sea ice and icebergs in the shelf sea environment adjacent to Antarctica. Credit: Mike Meredith, British Antarctic Survey

Scientists have made a new discovery that challenges previous understanding of the relationship between the polar Southern Ocean, next to Antarctica, and carbon dioxide levels in the atmosphere. Their findings show that, contrary to existing assumptions, biological processes far out at sea are the most important factors determining how the ocean absorbs carbon dioxide. The results are published this week in the journal *Science Advances*.

Carbon dioxide is absorbed in the surface ocean and stored in the deep seas over a timescale of 100s to 1000s years. The Southern Ocean plays a critical role in how this [carbon dioxide](#) is taken out of the atmosphere and knowing how it functions helps scientists understand dramatic climate transitions in the past, such as the ice ages, and better predict future climate change. It is commonly thought that the transformation of the water from light to dense—caused by cooling at the ocean's surface—is crucial in determining whether carbon is released to the atmosphere or trapped in the deep ocean. As a result, current research is often focused on the shallow seas right next to the Antarctic continent, where most of this transformation takes place.

The team, led by the University of Southampton with British Antarctic Survey, University of East Anglia and the Alfred Wegener Institute in Germany, studied the ocean circulation and carbon concentration of the Weddell Gyre, a region of critical importance for carbon removal from the atmosphere lying east of the Antarctic Peninsula. They studied data collected as part of the ANDREX project (Antarctic Deep water Rates of Export) which measured the physical, biological and chemical properties of the water of the waters in the gyre between 2008 and 2010.



The ANDREX project measures the properties of the Weddell Gyre, east of the Antarctic Peninsula. Credit: Mike Meredith, British Antarctic Survey

By studying this data, the team showed that the dominant factor driving the uptake of carbon from atmosphere to ocean is not related to dense water formation in the shallow seas immediately next to Antarctica, but rather to [biological processes](#) further out to sea. As phytoplankton in the center of the gyre grow then sink, they remove carbon from the surface

of the ocean, causing an uptake of carbon dioxide from the atmosphere—a process known as the "biological carbon pump." The data considered in this study showed unambiguously that, in the Weddell Gyre, this is the dominant process enabling the uptake of carbon dioxide from the atmosphere and its removal to the deep ocean.

Graeme MacGilchrist, who led the study for the University of Southampton said: "The results carry implications for our understanding of how the high-latitude Southern Ocean, close to the Antarctic continent, influences atmospheric carbon and global climate on 100 to 1000-year timescales. This is important both for our understanding of climate transitions in the past, such as the ice ages, as well as our projections of future climate change. We also expect that it will help to shift the focus of future research towards the critical processes taking place in the Antarctic Gyres, rather than the historical focus on the shelf-sea regions."



Collecting water samples from the deep ocean. Credit: Sinhue Torres Valdes, Alfred Wegener Institute

Michael Meredith from British Antarctic Survey said: "The Southern Ocean is a hugely important region for the drawdown of [carbon](#) dioxide from the [atmosphere](#), with major impacts on global climate. This study overturns a commonly-held belief relating to how this works, and emphasizes the need for joined-up biological and physical studies in the open-[ocean](#) regions some distance offshore from the Antarctic continent. This will be a key priority going forward, in order to improve our ability to reliably predict future climates."

More information: G.A. MacGilchrist et al., "Reframing the carbon cycle of the subpolar Southern Ocean," *Science Advances* (2019). DOI: 10.1126/sciadv.aav6410 , advances.sciencemag.org/content/5/8/eaav6410

Provided by University of Southampton

Citation: Research sheds new light on Antarctic control of global climate (2019, August 28) retrieved 19 April 2024 from <https://phys.org/news/2019-08-antarctic-global-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.