

## Astronomers reveal true colors of evolving galactic beasts

August 7 2019



Figure 1 - Red quasar: A brief transitional phase where the young quasar is enshrouded in gas and dust. This phase is potentially associated with young jets and strong winds, which ultimately drive away the obscuring dust. Credit: S. Munro



Astronomers have identified a rare moment in the life of some of the universe's most energetic objects. Quasars were first observed 60 years ago, but their origins still remain a mystery. Now researchers at Durham University, UK, have spotted what they suggest is a "brief transition phase" in the development of these galactic giants that could shed light on how quasars and their host galaxies evolve. The new research is published in *Monthly Notices of the Royal Astronomical Society*.

Quasars are powered by the energy from <u>supermassive black holes</u> at their centers as they feed on surrounding gases. They are thousands of times brighter than galaxies like our Milky Way and the majority are blue in color. However, a significant number are red as they are viewed through huge clouds of dust and gas that obscure them from view.

The conventional view of red quasars is that they are actually blue quasars that are angled away from our line-of-sight. Instead, the Durham team has ruled this model out and have shown that red quasars are likely to be the result of a brief, but violent, <u>phase</u> in the evolution of <u>galaxies</u> when the black hole ejects a large amount of energy into the surrounding clouds of dust and gas. This injection of energy blows away the dust and gas to reveal a blue quasar.

Observations using <u>radio telescopes</u> support this theory by showing that black holes at the center of red quasars produce a greater amount of radio emission than those at the center of blue quasars.

Lead author Lizelke Klindt, a Ph.D. researcher in Durham University's center for Extragalactic Astronomy, said: "How quasars develop has been the cause of significant uncertainty.





Figure 2 - Blue quasar: An unobscured quasar with signatures of evolved jets and less extreme winds that ultimately shut down star formation. Credit: S. Munro

"What our results suggest is that quasars undergo a brief transition phase, changing colour from red to blue, when they emerge from the deep shroud of dust and gas surrounding them.

"What we believe we are seeing is a rare but important step in the life of these galactic beasts during galaxy evolution when their <u>black holes</u> are starting to shape their environments."





Figure 3—red vs blue quasar: Red quasar: A brief transitional phase where the nascent quasar is enshrouded in gas and dust. This phase appears to be associated with young jets and strong winds, which ultimately drive away the obscuring gas and dust. Blue quasar: An unobscured quasar with signatures of evolved jets and less extreme winds. Credit: S. Munro

The researchers studied 10,000 red and blue <u>quasars</u> as they would have been seen seven to 11 billion years ago when the universe was relatively young using archival data from the Sloan Digital Sky Survey and the Very Large Array radio astronomy observatory.

They say their research could also tell us more about galaxy evolution. Co-author Professor David Alexander, Head of Astronomy, at Durham University, said: "We expect that during this transition phase the energy from the supermassive black hole will burn off the gas needed to form



stars.



Red and blue quasars are related within an evolutionary sequence that connects dust-obscured star formation with quasar activity through gas inflow via merging galaxies and outflows from the quasar. With this model the rare red quasar population represents a brief transitional phase between the starburst and the blue quasar phase during which winds and/or jets drive away the obscuring dust, revealing an unobscured blue quasar, and ultimately shutting down the star formation to form a dormant early-type galaxy. Credit: Gemini Observatory, GMOS-South, NSF; https://www.pxwall.com/4k-high-definition-galaxy-wallpaper/; Adapted by S. Munro

"Without the gas the galaxy cannot continue to grow, so what we are possibly seeing is the start of a quasar effectively ending the life of the galaxy by destroying the very thing it needs to survive."

The researchers say the next step in their research is to use more indepth data to understand the finer details of this transition phase.

**More information:** L Klindt et al. Fundamental differences in the radio properties of red and blue quasars: evolution strongly favoured over orientation, *Monthly Notices of the Royal Astronomical Society* 



## (2019). DOI: 10.1093/mnras/stz1771

## Provided by Royal Astronomical Society

Citation: Astronomers reveal true colors of evolving galactic beasts (2019, August 7) retrieved 5 May 2024 from <u>https://phys.org/news/2019-08-astronomers-reveal-true-evolving-galactic.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.