


# Atomic 'Trojan horse' could inspire new generation of X-ray lasers and particle colliders

August 13 2019

---



Illustration, based on simulations, of the Trojan horse technique for the production of high-energy electron beams. A laser beam (red, at left) strips electrons (blue dots) off of helium atoms. Some of the freed electrons (red dots) get accelerated inside a plasma bubble (white elliptical shape) created by an electron beam (green). Credit: Thomas Heinemann/University of Strathclyde

How do researchers explore nature on its most fundamental level? They build "supermicroscopes" that can resolve atomic and subatomic details. This won't work with visible light, but they can probe the tiniest dimensions of matter with beams of electrons, either by using them directly in particle colliders or by converting their energy into bright X-rays in X-ray lasers. At the heart of such scientific discovery machines are particle accelerators that first generate electrons at a source and then boost their energy in a series of accelerator cavities.

Now, an international team of researchers, including scientists from the Department of Energy's SLAC National Accelerator Laboratory, has demonstrated a potentially much brighter electron source based on plasma that could be used in more compact, more powerful [particle accelerators](#).

The method, in which the electrons for the [beam](#) are released from neutral atoms inside the plasma, is referred to as the Trojan horse technique because it's reminiscent of the way the ancient Greeks are said to have invaded the city of Troy by hiding their forceful soldiers (electrons) inside a wooden horse (plasma), which was then pulled into the city (accelerator).

"Our experiment shows for the first time that the Trojan horse method actually works," says Bernhard Hidding from the University of Strathclyde in Glasgow, Scotland, the principal investigator of a study published today in *Nature Physics*. "It's one of the most promising methods for future electron sources and could push the boundaries of today's technology."

## **Replacing metal with plasma**

In current state-of-the-art accelerators, electrons are generated by shining laser light onto a metallic photocathode, which kicks electrons

out of the metal. These electrons are then accelerated inside metal cavities, where they draw more and more energy from a radiofrequency field, resulting in a high-energy electron beam. In X-ray lasers, such as SLAC's Linac Coherent Light Source (LCLS), the beam drives the production of extremely bright X-ray light.

But metal cavities can only support a limited energy gain over a given distance, or acceleration gradient, before breaking down, and therefore accelerators for high-energy beams become very large and expensive. In recent years, scientists at SLAC and elsewhere have looked into ways to make accelerators more compact. They demonstrated, for example, that they can replace metal cavities with plasma that allows much higher acceleration gradients, potentially shrinking the length of future accelerators 100 to 1,000 times.

The new paper expands the plasma concept to the electron source of an accelerator.

"We've previously shown that plasma acceleration can be extremely powerful and efficient, but we haven't been able yet to produce beams with high enough quality for future applications," says co-author Mark Hogan from SLAC. "Improving beam quality is a top priority for the next years, and developing new types of electron sources is an important part of that."


According to [previous calculations](#) by Hidding and colleagues, the Trojan horse technique could make electron beams 100 to 10,000 times brighter than today's most powerful beams. Brighter electron beams would also make future X-ray lasers brighter and further enhance their scientific capabilities.

"If we're able to marry the two major thrusts—high acceleration gradients in plasma and beam creation in plasma—we could be able to

build X-ray lasers that unfold the same power over a distance of a few meters rather than kilometers," says co-author James Rosenzweig, the principal investigator for the Trojan horse project at the University of California, Los Angeles.

## Producing superior electron beams

The researchers carried out their experiment at SLAC's Facility for Advanced Accelerator Experimental Tests (FACET). The facility, which is currently undergoing a major upgrade, generates pulses of highly energetic electrons for research on next-generation accelerator technologies, including plasma acceleration.



An electron bunch from SLAC's FACET facility (bright spot at right) passes through hydrogen plasma (purple), which creates a plasma bubble (blue). As the bubble moves through the plasma at nearly the speed of light, a laser pulse strips electrons (white dots) off of neutral helium atoms inside the plasma. The

released electrons are trapped in the tail of the bubble where they gain energy (bright spot at left). Credit: Greg Stewart/SLAC National Accelerator Laboratory

First, the team flashed [laser light](#) into a mixture of hydrogen and helium gas. The light had just enough energy to strip electrons off hydrogen, turning neutral hydrogen into plasma. It wasn't energetic enough to do the same with helium, though, whose electrons are more tightly bound than those for hydrogen, so it stayed neutral inside the plasma.

Then, the scientists sent one of FACET's electron bunches through the plasma, where it produced a plasma wake, much like a motorboat creates a wake when it glides through the water. Trailing electrons can "surf" the wake and gain tremendous amounts of energy.

## **More R&D work ahead**

But before applications like compact X-ray lasers could become a reality, much more research needs to be done.

Next, the researchers want to improve the quality and stability of their beam and work on better diagnostics that will allow them to measure the actual beam brightness, instead of estimating it.

These developments will be done once the FACET upgrade, FACET-II, is completed. "The experiment relies on the ability to use a strong electron beam to produce the plasma wake," says Vitaly Yakimenko, director of SLAC's FACET Division. "FACET-II will be the only place in the world that will produce such beams with high enough intensity and energy."



In this study, the trailing electrons came from within the plasma (see animation above and movie below). Just when the electron bunch and its wake passed by, the researchers zapped the helium in the plasma with a second, tightly focused laser flash. This time the light pulse had enough energy to kick electrons out of the helium atoms, and the electrons were then accelerated in the wake.

The synchronization between the electron bunch, rushing through the plasma with nearly the speed of light, and the laser flash, lasting merely a few millionths of a billionth of a second, was particularly important and challenging, says UCLA's Aihua Deng, one of the study's lead authors: "If the flash comes too early, the electrons it produces will disturb the formation of the plasma wake. If it comes too late, the [plasma](#) wake has moved on and the electrons won't get accelerated."

The researchers estimate that the brightness of the electron beam obtained with the Trojan horse method can already compete with the brightness of existing state-of-the-art electron sources.

"What makes our technique transformative is the way the electrons are produced," says Oliver Karger, the other lead author, who was at the University of Hamburg, Germany, at the time of the study. When the electrons are stripped off the helium, they get rapidly accelerated in the forward direction, which keeps the beam narrowly bundled and is a prerequisite for brighter beams.

**More information:** A. Deng et al. Generation and acceleration of electron bunches from a plasma photocathode, *Nature Physics* (2019). [DOI: 10.1038/s41567-019-0610-9](https://doi.org/10.1038/s41567-019-0610-9)

Provided by SLAC National Accelerator Laboratory

Citation: Atomic 'Trojan horse' could inspire new generation of X-ray lasers and particle colliders (2019, August 13) retrieved 19 April 2024 from <https://phys.org/news/2019-08-atomic-trojan-horse-x-ray-lasers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.