

# Climate change could shrink oyster habitat in California

August 6 2019, by Kat Kerlin

---



A boulder of native oysters near Tomales Bay, California. Credit: UC Davis

Ocean acidification is bad news for shellfish, making it harder for them to form their calcium-based shells. But several other factors related to climate change could also make California bays less hospitable to shelled organisms like oysters, which are a key part of the food web.

Changes to water temperature and chemistry resulting from human-caused [climate change](#) could shrink the prime habitat and farming locations for oysters in California bays, according to a new study from the University of California, Davis.

The study, published today in the journal *Limnology and Oceanography*, shows that changes to dissolved [oxygen levels](#), [water temperature](#), and salinity could have an even greater impact than [ocean acidification](#) on oyster growth in estuaries and bays.

"The study demonstrates that focusing on [ocean](#) acidification alone is misguided," said UC Davis Professor Ted Grosholz, who led the study with funding from California Sea Grant. "Many climate-related stressors contribute to the projected shrinkage of the estuarine zone, where oysters and likely other shellfish would need to flourish."

## **What's different about estuaries**

California is home to two oyster species, which make their homes in partially enclosed estuaries and bays like Tomales Bay, where Grosholz and colleagues focused their research. The native Olympia oyster provides essential structure and food for other species. The non-native Pacific oyster is commonly farmed for food in the state, supporting a \$25-million industry.

The growing problem of ocean acidification—caused by increased [carbon dioxide](#) (CO<sub>2</sub>) in the atmosphere—has been a subject of concern among oyster growers and biologists in California.

In the estuaries and bays where oysters grow, ocean acidification is a much more complicated process than in the open ocean. Out at sea, atmospheric carbon drives the process of ocean acidification—as concentrations go up in the atmosphere, more carbon dioxide dissolves into the water, making it more acidic.

What's different about estuaries, said Grosholz, is there are many other inputs that can influence acidity.

"In an estuary, you have freshwater coming down from rivers, and there are plants, macroalgae, and phytoplankton that are much more abundant and greatly influence pH," he said.

This leads to daily cycle of changes in pH (the measure of acidity) that can far exceed pH changes in the ocean.



Professor Ted Grosholz pilots the boat while UC Davis graduate student Jordan Hollarsmith (left) and James Farlin (far right) of the UC Davis Bodega Marine Laboratory prepare to dive in California's Tomales Bay. Credit: Manon Picard/UC Davis

Estuarine organisms have evolved with this variable pH regime, which could make them more resilient to ocean acidification. But other factors important to oyster health, such as temperature, salinity and dissolved oxygen, are also projected to change with climate change.

## Exploring impacts

Grosholz started the research project in 2014 with UC Davis researcher Ann Russell to explore the impacts of acidification in estuaries and tease out other potential climate change impacts on oyster growth and health.

From 2014 to 2017, the researchers planted juvenile oysters of both species in test beds at nine different locations in Tomales Bay. They monitored their health over one-month periods, during a variety of seasonal conditions. At the same time, they closely monitored water chemistry, temperature, and other conditions at each location, and measured the growth and mortality of the oysters.

They also conducted field experiments with oyster larvae to see how variations in conditions affected survival and settlement—the process by which shellfish change from free-floating larvae and settle on a hard surface.

## **A stressful combination of threats**

The study found that the lowest oyster growth and survival rates happened in the parts of the bay most impacted by river runoff and inflow of upwelled ocean waters. While oyster growth and survival rates improved in the spring and summer, upwelled ocean waters low in dissolved oxygen and pH remained detrimental to the oysters.

In California, climate change is expected to lead to increased variability in precipitation, higher water temperatures, and increased upwelling. The study suggests that this combination of effects would lead to greater stress on oysters, particularly at the edges of bays that connect to rivers and the ocean.

"This means the oysters will be getting squeezed at both sides, reducing the zone of habitat they can thrive in," said Grosholz.

The study provides insight for oyster restoration projects as well as commercial oyster farmers. For example, projections for suitable [oyster](#) habitat could help determine where to site projects and farms for the best chance of success.

"Oysters settle to rocks so they cannot move and track changing water conditions like some other organisms," said UC Davis graduate student Jordan Hollarsmith, who is first author of the study. "The goldilocks zones within California estuaries, where conditions are just right for maximum growth and survival, are incredibly important for the economic success of farmed oysters and the ecological success of native oysters. Our findings help us to understand the environmental factors that make these zones so favorable, which will aid in restoration and aquaculture efforts."

Provided by UC Davis

Citation: Climate change could shrink oyster habitat in California (2019, August 6) retrieved 20 April 2024 from <https://phys.org/news/2019-08-climate-oyster-habitat-california.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.