


Japanese trees synchronize allergic pollen release over immense distances

August 27 2019



Created from public open data by Ministry of the Environment pollen observation system "Hanako-san". The in-phase matrix visualized the annual changes of phase-synchronizations across the Japanese islands. Credit: Kenshi

Sakai, TUAT

Complaints of allergic rhinitis (hayfever) are common worldwide, affecting around 17 percent of the Japanese population in spring and summer (around 20 million people). In Japan, the main tree species causing hayfever are Japanese cedar and Japanese cypress, with a combined land area of over 7 million hectares. Their pollen is dispersed between February and May and causes a range of symptoms from itchy eyes and runny noses through to severe respiratory disorders.

Management of these symptoms is dependent on timely and accurate pollen forecasting. It is known that the pollen from these trees has alternate annual cycles of ON (abundant pollen) and OFF (lower amounts) but it is not yet known how the dispersal of pollen synchronizes across Japan as studies to date have been based on only time-limited and local data.

In a study published on August 7 in *Scientific Reports*, researchers from Tokyo University of Agriculture and Technology (TUAT) made use of publicly available data from the Japanese Ministry of the Environment's "Hanako-san" pollen observation system to investigate annual fluctuations in pollen distribution at 120 locations for 15 years and to clarify the spatial synchronization across the Japanese islands.

This research is the first of its kind to study this synchronicity at the national scale and operates across a wide range of academic fields such as nonlinear physics, ecology, and [environmental science](#). "We used a new 'in-phase and out-of-phase' approach as an update to the previous correlation-based methods," says co-author of the study Kenshi Sakai. "We found that the phase synchronizations within and between regions were strong in eastern regions of Japan, such as Hokkaido and Kanto,

but the synchronization in western regions was much weaker."

Remarkably, all 120 sites in Japan behaved perfectly in-phase in 2009-2010, even over distances of over 1600 km. In a dramatic contrast, almost perfect desynchronization occurred in 2015-2016. "We still don't know what causes these [regional differences](#) in the pollen dispersal behavior," says lead author Akira Ishibashi, "It could be related to correlations in environmental variabilities at the locations of the different tree populations, known as the Moran effect."

Further research will be needed to pinpoint the specific mechanisms behind the observed synchronicity, but the findings will be useful for application in numerical forecasting of pollen abundance, which is crucial to clinicians and individuals suffering with hayfever.

"These findings and our proposed technique with an in-phase matrix have other applications outside of improving the prediction accuracy of allergenic [pollen](#)," says Sakai. "The new technique could also be used to quantitatively explain other environmental changes throughout Japan."

More information: Akira Ishibashi et al, Dispersal of allergenic pollen from *Cryptomeria japonica* and *Chamaecyparis obtusa*: characteristic annual fluctuation patterns caused by intermittent phase synchronisations, *Scientific Reports* (2019). [DOI: 10.1038/s41598-019-47870-6](#)

Provided by Tokyo University of Agriculture and Technology

Citation: Japanese trees synchronize allergic pollen release over immense distances (2019, August 27) retrieved 30 May 2023 from <https://phys.org/news/2019-08-japanese-trees-synchronize-allergic-pollen.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.