

Study finds native bighorn sheep herds retain migratory diversity

August 2 2019, by Rachel Hergett



A big horn ram looks back as yearling big horn sheep eat hay off a snow field while Montana State University students and faculty in the Department of Ecology, volunteer with Montana Fish, Wildlife and Parks officials and biologists in capturing and relocating big horn sheep on Saturday, Jan. 20, 2018, in the Quake Lake area, south of Ennis, as part of MSU research on repopulating historic big horn sheep grounds and landscape. Credit: MSU, Adrian Sanchez-Gonzalez



On the surface, bighorn sheep migration is like that of many other large mammals, moving to higher elevations as snow melts in the springtime then returning to lower ground to forage as winter sets in.

But a study published this month by Montana State University researchers has delved deeper, finding diverse patterns of migration among different sheep populations. Understanding these patterns could one day help wildlife officials make more informed management decisions.

The paper, "Characterizing <u>population</u> and individual <u>migration patterns</u> among native and restored <u>bighorn sheep</u> (Ovis Canadensis)," was published in the journal *Ecology and Evolution* in July.

"This paper is the first to quantify migratory diversity in ungulates and start to ask questions about how our understanding of migratory diversity can influence sheep populations and their restoration," said lead author Blake Lowrey, who is now a postdoctoral researcher in MSU's Department of Ecology in the College of Letters and Science. Lowrey completed his Ph.D. in November.

The paper draws from an extensive data set compiled by the Greater Yellowstone Mountain Ungulate Project and the Montana Bighorn Initiative, both of which are led by MSU ecology professor and coauthor Robert Garrott. Data was collected from 209 female bighorn sheep in the Rocky Mountains in collaboration with national parks, state wildlife agencies, land managers and researchers in Montana, Wyoming, Idaho and Colorado. Rather than focus on a single herd, the partnership allows for comparative studies across 18 populations of bighorn sheep.

Between 2008 and 2017, researchers fit bighorn sheep with GPS collars to track their movements. Blood was drawn to obtain measures of health in each animal. Nasal and tonsil swabs provided insight for disease and



pathogen work. According to Garrott, it's the largest coordinated research effort on bighorn sheep ever undertaken.

"Putting that all together provides a really rich data set to address a number of important topics related to bighorn sheep management," Garrott said.

Lowrey, a spatial ecologist who studies distribution of animal species over the landscape, used the locations collected from the GPS collars to investigate migration patterns in the bighorn sheep. He defined core summer and winter ranges for individuals and measured elevation and geographic distances between the ranges. These migration patterns were then compared between types of herds: native herds that have never been removed from their historic ranges and herds within the historic bighorn sheep range in the Rockies but had been restored or augmented through species translocation.

"Migratory patterns in ungulates are thought to be learned," Lowrey said. "If you remove animals from the landscape, you lose the herd memory needed to maintain a diversity of migratory patterns."

In his research, Lowrey found not only do the bighorn sheep in native populations tend to migrate more on average, they also exhibit longer migrations and have more diversity in how they migrate. Sheep within a single native herd may migrate upward of 25 miles, they may travel short distances between high and low elevations, or they may stay at high elevations year-round. Yet within the restored and augmented populations, there is less variety. Most individuals have the same relatively short migration or do not migrate at all.

"Blake's work is the first that has really demonstrated this contrast in migratory diversity between restored and <u>native populations</u>," Garrott said.



While researchers have yet to determine what effect these patterns may have on bighorn sheep populations, Lowrey notes that diversity has important benefits in migratory fish and bird populations.

"Practically, it's not putting all of your eggs in one basket," he said. "Without migratory diversity, an extreme winter event or other negative factor could negatively impact all of the population. Migratory diversity may buffer bighorn sheep and not expose all of the population to the same harmful conditions."

As part of his dissertation, Lowrey published papers on mountain goat habitat and range expansion, disease in mountain goats, and the niche overlap between native bighorn sheep and introduced mountain goats. And he said there's more coming. Lowrey also plans to look at the effect of landscape characteristics such as spring green-up, road density and other development as factors that may influence migratory patterns of bighorn <u>sheep</u>.

"The work that he's doing is giving us a stronger knowledge of how animals use the landscape and what they might need," Garrott said. "This knowledge should help inform restoration and management in the future."

More information: Blake Lowrey et al, Characterizing population and individual migration patterns among native and restored bighorn sheep (Ovis canadensis), *Ecology and Evolution* (2019). <u>DOI:</u> <u>10.1002/ece3.5435</u>

Provided by Montana State University

Citation: Study finds native bighorn sheep herds retain migratory diversity (2019, August 2)



retrieved 2 May 2024 from <u>https://phys.org/news/2019-08-native-bighorn-sheep-herds-retain.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.