

## Physicists solve 2,000-year-old optical problem

August 9 2019, by Bob Yirka



(a) Geometry of the problem and notation used for the distances. The origin of the coordinate system is located at the center of the input surface z a 0, 0??
(b) Zoom showing the notation for the unit vectors.

A trio of physicists from the National Autonomous University of Mexico and Tec de Monterrey has solved a 2,000-year-old optical problem—the Wasserman-Wolf problem. In their paper published in the journal *Applied Optics*, Rafael González-Acuña, Héctor Chaparro-Romo, and Julio Gutiérrez-Vega outline the math involved in solving the puzzle, give some examples of possible applications, and describe the efficiency of the results when tested.

Over 2,000 years ago, Greek scientist Diocles recognized a problem with <u>optical lenses</u>—when looking through devices equipped with them, the edges appeared fuzzier than the center. In his writings, he proposed that the effect occurs because the lenses were spherical—light striking at an angle could not be focused because of differences in refraction. Isaac



Newton was reportedly stumped in his efforts to solve the problem (which became known as <u>spherical aberration</u>), as was Gottfried Leibniz.

In 1949, Wasserman and Wolf devised an analytical means for describing the problem, and gave it an official name—the Wasserman-Wolf problem. They suggested that the <u>best approach</u> to solving the problem would be to use two aspheric adjacent surfaces to correct aberrations. Since that time, researchers and engineers have come up with a variety of ways to fix the problem in specific applications—most particularly cameras and telescopes. Most such efforts have involved creating aspherical lenses to counteract refraction problems. And while they have resulted in improvement, the solutions have generally been expensive and inadequate for some applications.

Now, a means for fixing the problem with any size <u>lens</u> has been found by González-Acuña, Chaparro-Romo and Gutiérrez-Vega, described in a lengthy math formula. It is based on describing ways in which the shape of a second aspherical surface needs to be given a first surface, along with object-image distance. In essence, it relies on a second surface fixing problems with the first surface. The result is elimination of spherical aberration.

**More information:** Rafael G. González-Acuña et al. General formula to design a freeform singlet free of spherical aberration and astigmatism, *Applied Optics* (2019). DOI: 10.1364/AO.58.001010



## © 2019 Science X Network

Citation: Physicists solve 2,000-year-old optical problem (2019, August 9) retrieved 2 May 2024 from <u>https://phys.org/news/2019-08-physicists-year-old-optical-problem.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.