


# Ring-shaped multi-carbon compound cyclocarbon synthesized

August 16 2019, by Bob Yirka




From left to right, precursor molecule  $\text{C}_{24}\text{O}_6$ , intermediates  $\text{C}_{22}\text{O}_4$  and  $\text{C}_{20}\text{O}_2$  and the final product cyclo[18]carbon  $\text{C}_{18}$  created on surface by dissociating CO masking groups using atom manipulation. The bottom row shows atomic force microscopy (AFM) data using a CO functionalized tip. Credit: IBM Research

A team of researchers from Oxford University and IBM Research has for the first time successfully synthesized the ring-shaped multi-carbon compound cyclocarbon. In their paper published in the journal *Science*, the group describes the process they used and what they learned about


the bonds that hold a cyclocarbon together.

Carbon is one of the most abundant elements, and has been found to exist in many forms, including diamonds and graphene. The researchers with this new effort note that much research has been conducted into the more familiar forms (allotropes) how they are bonded. They further note that less well-known types of [carbon](#) have not received nearly as much attention. One of these, called cyclocarbon, has even been the topic of debate. Are the two-neighbor forms bonded by the same length bonds, or are there alternating bonds of shorter and longer lengths? The answer to this question has been difficult to find due to the high reactivity of such forms. The researchers with this new effort set themselves the task of finding the answer once and for all.


The team's approach involved creating a precursor molecule and then whittling it down to the desired form. To that end, they used [atomic force microscopy](#) to create linear lines of [carbon atoms](#) atop a copper substrate that was covered with salt to prevent the carbon atoms from bonding with the subsurface. They then joined the lines of atoms to form the carbon oxide precursor  $C_{24}O_6$ , a triangle-shaped form. Next, the team applied [high voltage](#) through the AFM to shear off one of the corners of the triangle, resulting in a  $C_{22}O_4$  form. They then did the same with the other two corners. The result was a  $C_{18}$  ring—an 18-atom cyclocarbon. After creating the ring, the researchers found that the bonds holding it together were the alternating long- and short-type bonds that had been previously suggested.



Cyclo[18]carbon structure model. Credit: IBM Research



Cyclo[18]carbon. Bottom: artistic 3-dimensional representation of AFM data (at increased tip-sample distance); top: molecular structure model. Credit: IBM Research



Cyclo[18]carbon. Bottom: artistic 3-dimensional representation of AFM data (at small tip-sample distance); top: molecular structure model. Credit: IBM Research

The researchers note that their technique allows for further experimentation—they could try fusing cyclocarbons, for example, to create new kinds of molecules and, by extension, materials. They further note that the technique also opens the door to the creation of new types of electronics based on single-electron transfer.

**More information:** Katharina Kaiser et al. An sp-hybridized molecular carbon allotrope, cyclo[18]carbon, *Science* (2019). [DOI: 10.1126/science.aay1914](https://doi.org/10.1126/science.aay1914)

IBM Research blog: [www.ibm.com/blogs/research/201... imaging-](https://www.ibm.com/blogs/research/2019-11-14-imaging-cyclo-18-carbon)

[cyclocarbon/](#)

© 2019 Science X Network

Citation: Ring-shaped multi-carbon compound cyclocarbon synthesized (2019, August 16)  
retrieved 3 May 2024 from

<https://phys.org/news/2019-08-ring-shaped-multi-carbon-compound-cyclocarbon.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--